




Research Article

Phytochemical Analysis, Acute Toxicity, as well as Antihyperglycemic and Antidiabetic Activities of *Corchorus olitorius* L. Leaf Extracts

Rebecca Nakaziba ^{1,2}, Aloysius Lubega ³, Jasper Ogwal-Okeng ², and Paul E. Alele ¹

¹Department of Pharmacology and Therapeutics, Mbarara University of Science and Technology, P.O Box, Mbarara 1410, Uganda

²Department of Pharmacology and Therapeutics, Lira University, P.O Box, Lira 1035, Uganda

³Department of Pharmacology and Therapeutics, Makerere University College of Health Sciences, P.O Box, Kampala 7072, Uganda

Correspondence should be addressed to Rebecca Nakaziba; d2rebecca1@gmail.com

Received 8 March 2022; Revised 15 June 2022; Accepted 1 July 2022; Published 18 July 2022

Academic Editor: Ghadir A. El-Chaghably

Copyright © 2022 Rebecca Nakaziba et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Background and Aim. Diabetes mellitus is a metabolic disorder that has no known cure with continuous endeavors to find a therapy for the condition. According to some studies, traditional leafy vegetables could prevent and manage diabetes by modifying the carbohydrate and lipid metabolism. In this study, a phytochemical analysis, acute toxicity, as well as antihyperglycemic and antidiabetic activity testing of the methanolic, diethyl ether, and aqueous leaf extracts of *Corchorus olitorius* L. was performed. **Materials and Methods.** Methanolic, diethyl ether, and aqueous leaf extracts of *Corchorus olitorius* L. were prepared by serial extraction. Phytochemical analysis was performed following standard methods. 52 mice were separated into 13 groups (A–M) of 4 and received extracts' doses ranging from 1000 mg/kg to 5000 mg/kg for the acute toxicity testing. For the antihyperglycemic and antidiabetic activities testing, 48 rats were divided into 8 groups of 6 and received 500 mg/kg of each extract. 10 mg/kg of glibenclamide and distilled water were used as controls. Data were analyzed using Prism GraphPad version 8.0.2 (263). **Results.** Phytochemical analysis revealed the presence of alkaloids, reducing sugars, saponins, and terpenoids. There were no acute toxicity signs observed in this study. *Corchorus olitorius* L. extracts demonstrated moderate antihyperglycemic and antidiabetic activities. The methanolic extract exhibited the highest degree of antihyperglycemic activity. However, there was no statistically significant difference between the extracts and the negative control ($p > 0.05$), but with glibenclamide ($p < 0.01$). **Conclusion.** *Corchorus olitorius* L. is a safe and potential postprandial antidiabetic vegetable that could minimize the rise in blood glucose after a meal. We therefore recommend further investigations into the antidiabetic properties of the vegetable using purified extracts.

1. Introduction

Corchorus olitorius L. (*C. olitorius*), also known as Tossa jute in English, is a traditional medicinal vegetable belonging to the genus *Corchorus* and the family of Tiliceae [1]. The vegetable is prevalent in northern Australia, tropical Africa, and India [1]. In these places, *C. olitorius* leaves commonly are served as a vegetable frequently in stews eaten with staple starchy foods [1]. *C. olitorius* is an annual herb growing to a height of 2.4 m [1]. It has alternating, finely indented margin leaves. *C. olitorius* has small yellow flowers comprising of five

petals that form a brown multiseeded pod later in life [1]. It is proliferated by seeds [2] much as it can be accepted as a wild vegetable in crop fields or cultivated in family gardens [2]. *C. olitorius* flourishes best throughout rainy seasons, nonetheless it is drought resilient [1, 2]. In West Africa, *C. olitorius* is used to treat malaria, typhoid fever, female infertility, heart failure, and ulcers [1, 3, 4]. In addition, it is used as a therapy for fevers, colds, constipation, and tumors [5]. The vegetable is also reported to have antiobesity [6, 7], anti-inflammatory [8, 9], antimicrobial [10, 11], and gastroprotective properties [12] along with antidiabetic effects [7, 13, 14].

Diabetes mellitus (DM), a disorder of mainly glucose metabolism, is one of the most common noncommunicable diseases (NCDs). In 2015, DM caused 5 million deaths and an estimated 12% of global expenditure [15–17]. The disease occurs in two types, that is, 1 and 2. Type 1 accounts for 2–10% of all diabetic cases in the world [18] and for >85% of diabetes registered amongst youth below 20 years of age [19]. It occurs as a result of autoimmune destruction of pancreatic beta cells with a subsequent total insulin deficiency [18]. Type 2 results from insensitivity to insulin and/or inadequate compensatory insulin secretion with over 90% of all diabetic cases in the world [15]. In Africa, diabetes occurred among 7.1% of the population in 2014, and the overall cost of DM was USD 19.45 million in 2015 [20]. More than 90% of diabetic cases in Africa are type 2 [20, 21]. In Uganda, the prevalence of diabetes was 1.9% in 2016 [22]. Diabetes is managed by dietary modification, exercise, and pharmacotherapy [23]. However, there is as yet no known cure for DM. This has driven the affected individuals' world over to opt for alternative medication including traditional herbal remedies, while a lot of research is being conducted to evaluate and document plants with glycemic effects [24–31]. In Uganda and other parts of the world, studies have revealed a number of plants that are used by communities in the management of diabetes [32, 33]. According to a study conducted by Momo et al [31], traditional leafy vegetables could prevent and manage diabetes by modifying the carbohydrate and lipid metabolism. *C. olitorius* has indicated a reduction in serum blood glucose levels in a few studies [34].

In the current study, we performed a phytochemical analysis, acute toxicity, as well as antihyperglycemic and antidiabetic activity testing of the methanolic, aqueous, and diethyl ether extracts of *C. olitorius* leaves in laboratory animals. This was done to confirm some of the reports about the vegetable as well as to assess its safety.

2. Materials and Methods

2.1. Laboratory Animals. Laboratory mice and rats from the animal house of the Department of Pharmacology and Therapeutics, Makerere College of Health Sciences, were used for this study. The animals (male and female) were housed in wooden cages at temperature between 25 and 28°C with a normal 12-hour/12-hour light/dark cycle for at least 2 weeks before the study. They were fed with standard rat feeds and water. Their weight ranged from 18 to 31 g and 128 g to 221 g for mice and rats, respectively. Pregnant and breast-feeding animals were excluded from the study.

2.2. Extract Preparation. The *C. olitorius* leaves were harvested from Olago village, Lira district (northern Uganda), and packaged in polyethylene bags. They were then transported to the Department of Pharmacology and Therapeutics, Makerere University College of Health Sciences, where they were washed clean and spread to air-dry inside the laboratory to a constant weight (900 g). The fully dried leaves were then ground to a coarse powder and extracted by means of solvents (diethyl ether, methanol, and distilled water) in a serial

extraction method as described by Das, Tiwari, and Shrivastava [35]. This was done so as to extract compounds of varied polarity. Briefly, 600 g of *C. olitorius* leaves powder were soaked in 3 L of diethyl ether (BDH AnalaR) in a flat bottomed conical flask and tightly closed while shaking every 6 hours for two days (48 hours.). The resultant solution was filtered using Whatman filter paper (No. 1), and the filtrate was concentrated by means of a rotary evaporator at 20–30°C. The remains were then spread in the laboratory and allowed to dry. After drying, it was soaked in 3 L of methanol (UltraPure Solutions, Inc. 11485, Commercial Parkway, Castroville, CA 95012), and the process was repeated as with diethyl ether. Last, the dry residue was soaked in 3 L of distilled water while shaking every 6 hours for two days (48 hours). The resultant solution was also filtered using Whatman filter paper (No. 1) and freeze dried. After evaporation of the solvents, the extracts were accordingly labeled and stored in a refrigerator at –80°C. A voucher specimen was kept at a herbarium at the Department of Botany, Makerere University, Kampala, Uganda (# 50906).

2.3. Phytochemical Analysis. The phytochemical analysis was performed as follows.

2.3.1. Tannins. 2 mL of 5% FeCl₃ were added to 2 mL of aqueous extract and observed for a yellow/brown precipitate formation [36].

2.3.2. Alkaloids. 1.5 mL of 1% HCl was added to 2 mL of the methanol extract filtrate, the solution was heated in a waterbath (5 minutes), and then 6 drops of Mayor's reagent was added and observed for an orange precipitate formation [37].

2.3.3. Saponins. 2 g powder of the aqueous extract was mixed with few drops of olive oil, shaken vigorously, and observed for formation of a stable persistent froth and an emulsion [38].

2.3.4. Cardiac Glycosides. To 2 mL methanol extract filtrate, 1 mL glacial acetic acid and 1-2 drops of FeCl₃ were added followed by 1 mL of concentrated H₂SO₄. Formation of a brown ring at the interface indicated the presence of deoxy-sugar [39, 40].

2.3.5. Terpenes. To 2 mL of the aqueous extract, 5 mL CHCl₃, 2 mL acetic anhydride, and a drop of concentrated H₂SO₄ were added carefully to form a layer. Formation of a reddish/brown coloration at the interface indicated the presence of terpenes [41].

2.3.6. Reducing Sugars. To 2.5 mL of Benedict's solution was added 0.5 g of the aqueous extract in a test tube. The mixture was warmed over a waterbath for about 5 minutes. Observation of green/red or yellow coloration indicated the presence of reducing sugars [42].

2.3.7. Nonreducing Sugars. 3 mL of the aqueous extract was mixed with 2 drops of dilute iodine solution and boiled for 5 minutes. Observation of blue color that disappeared upon boiling and reappeared upon cooling indicated the presence of nonreducing sugar [43].

2.4. Acute Toxicity Testing. The extracts were orally administered following the protocol laid down by the Organization for Economic Cooperation and Development [44], where the dose of the extract per animal body weight was calculated and administered to the animals suspended in distilled water. The animals were observed for mortality, behavioral changes, physical appearance, injury, and pain after 30 minutes, 1 hour, 2 hours, 4 hours, 8 hours, and once daily for 13 days. The control group was only given distilled water. Observations were recorded per group of animals. The experiment was performed as follows: a total of 52 mice (average weight 25 g) were divided into thirteen (13) groups (A–M) of four (4). Groups A–D, E–H, and I–L received 1000 mg/kg, 2000 mg/kg, 4000 mg/kg, and 5000 mg/kg of the diethyl ether, methanol, and aqueous extracts, respectively. Last, group M received 1 mL of distilled water.

2.5. The Antihyperglycemic and Antidiabetic Tests. This was performed as described by Mushtaq et al. [45] with minor modifications as follows. The rats were fasted overnight and randomly selected and distributed into eight groups ($n = 6$). To one of the groups, 1 mL of distilled water was orally administered. To three of the groups (for the antihyperglycemic test), the methanolic, diethyl ether, and aqueous extracts of *C. olitorius* (500 mg/kg body weight for each extract) were orally administered. After one hour, the animals were fed with glucose (2 g/kg body weight) [45]. To another set of three groups (for the antidiabetic test), alloxan (60 mg/kg; i.v) was given prior (3 days) to the study in order to induce diabetes. These too were given in 500 mg/kg of body weight of each of the extracts. The last group received glibenclamide (10 mg/kg body weight). A blood sample was then collected by pricking the tail vein of the rat using a scalp vein needle. Blood was gently milked, dropped on a glucose strip, and read using a glucometer (Code-free blood glucose test strips, Biosenser, Inc, Korea and Code free, Yeongtong-dong Yeontong-du, Suwoni-si, Kyonggi-go, Korea). Blood was withdrawn from the tail vein at 0, 30, 60, 90, 120, and 150 minutes of glucose administration for the antihyperglycemic test and the same time interval following test drug administration for the antidiabetic test. The glucose level is recorded in a table as indicated by the glucometer.

2.6. Data Management. Data was analyzed by Prism GraphPad version 8.0.2 (263) and expressed as mean + standard deviation. Significance level was set at $p \leq 0.05$ at 95% confidence interval (CI). The results are presented as tables and figures.

3. Results

3.1. Phytochemical Analysis. Phytochemical analysis revealed the presence of alkaloids, reducing sugars, and saponins among others (Table 1).

3.2. Acute Toxicity Testing. There were no observed acute toxicity effects of the *C. olitorius* extracts at the doses tested. Neither behavioral or appearance changes nor mortality was observed in all the groups. All the animals appeared and behaved normally (Tables 2 and 3).

3.3. Antihyperglycemic and Antidiabetic Activities Testing. To assess the antihyperglycemic activity of *C. olitorius*, the methanolic, diethyl ether, and aqueous extracts of *C. olitorius* leaves along with distilled water and glibenclamide were administered to rats an hour before glucose administration. The blood glucose level was then taken at different time intervals using a glucometer. For the antidiabetic test, the test animals were given alloxan three days before the experiment. *C. olitorius* extracts moderately exhibited antihyperglycemic and antidiabetic effects, although not comparable to glibenclamide (Table 4 and Figure 1).

P value < 0.05 was compared with positive control but > 0.05 with negative control for the antihyperglycemic test; P value = < 0.0001 was compared with water in the antidiabetic test (unpaired t -test with Welch's correction) for all the extracts. The overall P value calculation using the one-way ANOVA and multiple t -test was below 0.05 showing a statistically significant difference between the *C. olitorius* extracts and glibenclamide, whereas it was above 0.05 in comparison with water indicating no significant difference. However, the multiple t -test per time interval revealed a statistically significant difference between *C. olitorius* extracts and water at 30 minutes of glucose administration (P value < 0.05).

4. Discussion

Phytochemical analysis revealed the presence of alkaloids, glycosides, saponins, tannins, terpenoids, and reducing sugars (Table 1). This finding was consistent with a related study in India [46] and Bangladesh [47] but not in West Africa [48]. In another related study, alkaloids, tannins, and resins were found [34]. The discrepancy in phytochemical composition could be attributed to the differences in the climate of the regions. In addition, there was no observed acute toxicity effect of *C. olitorius* extracts seen in the current study (Table 3). This finding is similar to a set of related studies in Bangladesh and Nigeria where there were no observed acute toxicity effects in doses comparable to those of this study [47, 49]. In another related study, *C. olitorius* did not show any toxicity using enzyme markers in alloxan-induced diabetic rats [34]. This finding also supports the results of the present study. This implies that the vegetable is safe for consumption. To assess the antihyperglycemic activity of *C. olitorius* leaf extracts, the controls and the

TABLE 1: Phytochemical analysis of *C. olitorius*.

Phytochemicals	Present (+)/absent (-)
Alkaloid	+
Glycoside	+
Reducing sugar	+
Nonreducing sugar	-
Saponins	+
Tannins	+
Terpenoids	+

TABLE 2: Extract doses given.

Group	Extract	Dose (mg/kg)
A	Methanol	1000
B	Methanol	2000
C	Methanol	4000
D	Methanol	5000
E	Diethyl ether	1000
F	Diethyl ether	2000
G	Diethyl ether	4000
H	Diethyl ether	5000
I	Aqueous	1000
J	Aqueous	2000
K	Aqueous	4000
L	Aqueous	5000
M	Distilled water	0

TABLE 3: Acute toxicity of *C. olitorius* extracts.

Groups	Dose	Mortality (#)	Behavioral change (#)	Change in physical appearance (#)	Expression of pain (#)
A, E, I	1000 mg/kg	0	0	0	0
B, F, J	2000 mg/kg	0	0	0	0
C, G, K	4000 mg/kg	0	0	0	0
D, H, L	5000 mg/kg	0	0	0	0
M	0 mg/kg	0	0	0	0

#Number.

TABLE 4: The mean values of the plasma glucose level per group with the standard deviations.

T	M1	DE1	A1	M2	DE2	A2	Gl	W
0	6.850 ± 0.797	5.967 ± 0.625	5.317 ± 0.688	6.160 ± 0.953	5.900 ± 0.490	5.920 ± 0.746	3.633 ± 1.098	6.538 ± 1.327
30	7.767 ± 0.441	11.467 ± 0.644	9.983 ± 1.057	7.600 ± 0.797	5.460 ± 0.783	6.820 ± 1.062	9.600 ± 1.534	17.383 ± 1.326
60	7.583 ± 1.186	7.033 ± 1.529	8.333 ± 1.537	7.360 ± 1.555	5.860 ± 1.106	5.560 ± 0.757	4.367 ± 0.413	7.567 ± 0.378
90	6.783 ± 1.091	6.483 ± 1.592	6.450 ± 0.907	6.940 ± 1.001	6.620 ± 0.746	7.720 ± 1.893	3.167 ± 1.303	6.217 ± 1.353
120	5.583 ± 0.542	6.617 ± 0.571	6.217 ± 1.530	7.220 ± 0.576	6.680 ± 0.847	7.760 ± 0.358	2.500 ± 1.081	5.683 ± 1.118
150	5.267 ± 0.799	5.467 ± 1.549	5.317 ± 0.854	6.440 ± 0.777	7.400 ± 0.812	6.440 ± 0.643	2.517 ± 1.300	5.733 ± 0.668

T, time in minutes; M1, DE1, and Aq1, methanol, diethyl ether, and aqueous for extracts antihyperglycemic test; M2, DE2, and Aq2, methanol, diethyl ether, and aqueous extracts for the antidiabetic test; Gl, glibenclamide; W, distilled water.

C. olitorius extracts were administered an hour before glucose ingestion. Glibenclamide registered the lowest blood glucose at 0 hour of glucose administration. This was partly in agreement with the findings of Maxwell et al. in Nigeria [14] who tested the antidiabetic potential of the ethanolic seed extract of *C. olitorius* and found glibenclamide to have a

higher blood glucose lowering potential than the *C. olitorius* extract in normal glycemic rats. Meanwhile, there was no significant difference between the blood glucose levels of the *C. olitorius* extracts in comparison with the negative control at 0 hour. However, the blood glucose level for the methanolic extract was slightly higher than that of the negative




FIGURE 1: Blood glucose levels for the antihyperglycemic test.

control at 0 hour (Table 4). This indicates that the extracts have a minimal effect on baseline blood glucose. In an oral glucose tolerance test (OGTT) performed by Maxwell et al. [14], the ethanolic *C. olitorius* seed extract demonstrated a significant decrease in the blood glucose level from 30 minutes to 2 hours ($P < 0.001$). Similarly, in the current study, the different *C. olitorius* extracts demonstrated significant antihyperglycemic activity at 30 minutes of glucose ingestion ($p < 0.01$). Nevertheless, the overall p value ($p > 0.05$) in comparison with the negative control for the entire test was not statistically significant. This could be attributed to the fact that they (Maxwell et al.) [14] used the *C. olitorius* seed extract, yet the current study used leaves. There is also a probability that *C. olitorius* seeds exhibit a higher antihyperglycemic potential than the leaves. Comparison between the *C. olitorius* extracts and glibenclamide revealed that glibenclamide had a higher antihyperglycemic activity ($p < 0.01$) except with the aqueous extract at 30 minutes ($p = 0.63$). This too was similar to the findings of Maxwell et al. [14]. Their glibenclamide group had the highest reduction in blood glucose level among the test animals. This could be justified by the fact that glibenclamide is an already purified compound unlike the extracts. In another study conducted by Arise et al. [13], to establish the antidiabetic properties of *C. olitorius* in alloxan-induced diabetic rats, the ethanolic leaf extract exhibited antidiabetic activity. This finding relates to the outcome of the current study, although a methanolic extract was used in the current study. In addition, according to a study that tested the hypoglycemic effect of *C. olitorius* in albino rats, the methanolic extract demonstrated hypoglycemic activity, thereby, supporting the findings of the current study [7]. In a related study conducted by Sanjida et al. in Bangladesh [47], *C. olitorius* showed a dose-dependent reduction in the blood glucose level in an oral glucose tolerance test (OGTT) among test animals. This as well is in agreement with the first one hour findings of the current study. Furthermore, a study was conducted by Oboh et al. [50] to identify the antidiabetic potential of *C. olitorius*, and the plant demonstrated

antidiabetic activity in type 2 diabetic rats. This outcome also is in agreement with the findings of the current study, although the current effect was minimal. All the extracts in the present study moderated the rise in plasma glucose an hour after oral ingestion of glucose by the test animals. The methanolic extract showed the greatest antihyperglycemic activity within the first one hour. However, after the first one hour, there was no significant difference between the plasma glucose level of the extracts and water signifying a short-lived antihyperglycemic activity of the extracts. They therefore cannot be considered for long-term control of high blood sugar. In addition, the antihyperglycemic and antidiabetic activities of the extracts were less when compared to those of glibenclamide. This is attributable to the fact that glibenclamide is a pure compound. The antihyperglycemic and antidiabetic activities of *C. olitorius* are ascribed to the presence of alkaloids, which are believed to induce glucose uptake by pancreatic beta cells [51].

5. Conclusion and Recommendation

C. olitorius L. has moderate antihyperglycemic and antidiabetic activities with no toxic effects. Further in-depth studies to investigate the antihyperglycemic, antidiabetic, pharmacodynamics, and the pharmacokinetic properties of *C. olitorius* L. leaf extracts as a potential nutraceutical amongst diabetic patients should be performed using purified extracts.

Data Availability

The data used to support this study are available from the corresponding author upon request.

Ethical Approval

Ethical approval to conduct the study was provided by the Research and Ethics Committee (REC) of Mbarara University of Science and Technology, the Uganda National Council for Science and Technology (UNCST-HS2589), and the concerned laboratory administrators.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

RN conceptualized and designed the study, collected and analyzed data, and drafted the manuscript. AL performed laboratory tests and reviewed the manuscript. PA and JO supervised and reviewed the manuscript.

Acknowledgments

This study was supported by Lira University and the Department of Pharmacology and Therapeutics, Makerere University College of Health Sciences.

References

- [1] M. Loumerem and A. Alercia, "Descriptors for jute (*Corchorus olitorius* L.)," *Genetic Resources and Crop Evolution*, vol. 63, no. 7, pp. 1103–1111, 2016.
- [2] T. Begum and D. Kumar, "Usefulness of morphological characteristics for DUS testing of jute (*Corchorus olitorius* L. and *C. capsularis* L.)," *Spanish Journal of Agricultural Research*, vol. 9, no. 2, p. 473, 2011.
- [3] H. O. Adebo, L. E. Ahoton, F. J. B. Quenum, H. Adoukonou-Sagbadja, D. O. Bello, and C. A. A. M. Chrysostome, "Ethnobotanical knowledge of jute (*Corchorus olitorius* L.) in Benin," *European Journal of Medicinal Plants*, vol. 26, pp. 1–11, 2018.
- [4] D. Nyadanu, R. Adu Amoah, A. O. Kwarteng et al., "Domestication of jute mallow (*Corchorus olitorius* L.): ethnobotany, production constraints and phenomics of local cultivars in Ghana," *Genetic Resources and Crop Evolution*, vol. 64, no. 6, pp. 1313–1329, 2017.
- [5] M. M. Islam, "Biochemistry, medicinal and food values of jute (*Corchorus capsularis* L. and *C. olitorius* L.) leaf: a review," *International Journal of Enhanced Research In Science Technology & Engineering*, vol. 2, no. 11, pp. 135–144, 2013.
- [6] L. Wang, M. Yamasaki, T. Katsube, X. Sun, Y. Yamasaki, and K. Shiwaku, "Antiobesity effect of polyphenolic compounds from molokheiya (*Corchorus olitorius* L.) leaves in LDL receptor-deficient mice," *European Journal of Nutrition*, vol. 50, no. 2, pp. 127–133, 2011.
- [7] A. I. Airaodion, J. D. Akinmolayan, E. O. Ogbuagu, E. O. Airaodion, U. Ogbuagu, and O. O. Awosanya, "Effect of methanolic extract of *Corchorus olitorius* Leaves on hypoglycemic and hypolipidaemic activities in albino rats," *Asian Plant Research Journal*, vol. 2, pp. 1–13, 2019.
- [8] H. Handoussa, R. Hanafi, I. Eddiasty et al., "Anti-inflammatory and cytotoxic activities of dietary phenolics isolated from *Corchorus olitorius* and *Vitis vinifera*," *Journal of Functional Foods*, vol. 5, no. 3, pp. 1204–1216, 2013.
- [9] Y.-Y. Yan, Y. W. Wang, S. L. Chen, S. R. Zhuang, and C. K. Wang, "Anti-inflammatory effects of phenolic crude extracts from five fractions of *Corchorus olitorius* L.," *Food Chemistry*, vol. 138, no. 2–3, pp. 1008–1014, 2013.
- [10] S. Ilhan, F. Savaroglu, and F. Colak, "Antibacterial and antifungal activity of *Corchorus olitorius* L.(Molokhia) extracts," *International Journal of Natural and Engineering Sciences*, vol. 1, no. 3, 2007.
- [11] Z. A. Zakaria, M. N. Somchit, H. Zaiton et al., "The in vitro antibacterial activity of *Corchorus olitorius* extracts," *International Journal of Pharmacology*, vol. 2, no. 2, pp. 213–215, 2006.
- [12] R. Nakaziba, "Traditional uses of *Corchorus olitorius* L in Oyam District, northern uganda: a cross-sectional ethnobotanical survey," 2020, https://www.researchgate.net/publication/340189765_Traditional_uses_of_Corchorus_olitorius_L_in_Oyam_District_Northern_Uganda_A_cross-sectional_ethnobotanical_survey.
- [13] A. R. Olusanya, B. S. Ifeoluwa, A. J. Aboyewa, and B. Khadijat, "Antidiabetic and safety properties of ethanolic leaf extract of *corchorus olitorius* in alloxan-induced diabetic rats," *Diabetes Food Plan*, vol. 6, p. 57, 2018.
- [14] M. O. Egua, E. U. Etuk, S. O. Belloc, and S. W. Hassan, "Antidiabetic activity of ethanolic seed extract of *Corchorus olitorius*," *International Journal of Sciences: Basic and Applied Research*, vol. 12, no. 1, pp. 8–21, 2013.
- [15] Y. Zheng, S. H. Ley, and F. B. Hu, "Global aetiology and epidemiology of type 2 diabetes mellitus and its complications," *Nature Reviews Endocrinology*, vol. 14, no. 2, pp. 88–98, 2018.
- [16] G. Roglic, "WHO Global report on diabetes: a summary," *International Journal of Noncommunicable Diseases*, vol. 1, no. 1, p. 3, 2016.
- [17] K. Ogurtsova, J. da Rocha Fernandes, Y. Huang et al., "IDF Diabetes Atlas: global estimates for the prevalence of diabetes for 2015 and 2040," *Diabetes Research and Clinical Practice*, vol. 128, pp. 40–50, 2017.
- [18] D. M. Maahs, N. A. West, J. M. Lawrence, and E. J. Mayer-Davis, "Epidemiology of type 1 diabetes," *Endocrinology and Metabolism Clinics of North America*, vol. 39, no. 3, pp. 481–497, 2010.
- [19] J. M. Lawrence, D. A. Standiford, B. Loots et al., "Prevalence and correlates of depressed mood among youth with diabetes: the SEARCH for Diabetes in Youth study," *Pediatrics*, vol. 117, no. 4, pp. 1348–1358, 2006.
- [20] R. Atun, J. I. Davies, E. A. M. Gale et al., "Diabetes in sub-Saharan Africa: from clinical care to health policy," *Lancet Diabetes & Endocrinology*, vol. 5, no. 8, pp. 622–667, 2017.
- [21] S. D. Pastakia, C. Pekny, S. Manyara, and L. Fischer, "Diabetes in sub-Saharan Africa – from policy to practice to progress: targeting the existing gaps for future care for diabetes," *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, vol. 10, pp. 247–263, 2017.
- [22] S. Bahendeka, R. Wesonga, G. Mutungi, J. Muwonge, S. Neema, and D. Guwatudde, "Prevalence and correlates of diabetes mellitus in Uganda: a population-based national survey," *Tropical Medicine and International Health*, vol. 21, no. 3, pp. 405–416, 2016.
- [23] S. Madhu and S. Srivastava, "Diabetes mellitus: diagnosis and management guidelines," *Journal International Medical Sciences Academy*, vol. 1, pp. 47–50, 2015.
- [24] P. Tripathi, R. Srivastava, A. Pandey, R. Pandey, and S. Goswami, "Alternative therapies useful in the management of diabetes: a systematic review," *Journal of Pharmacy and BioAllied Sciences*, vol. 3, no. 4, p. 504, 2011.
- [25] E. A. Ryan, M. E. Pick, and C. Marceau, "Use of alternative medicines in diabetes mellitus," *Diabetic Medicine*, vol. 18, no. 3, pp. 242–245, 2001.
- [26] K. Hjelm and E. Mufunda, "Zimbabwean diabetics' beliefs about health and illness: an interview study," *BMC International Health and Human Rights*, vol. 10, no. 1, p. 7, 2010.
- [27] T. Salihu Shinkafi, L. Bello, S. Wara Hassan, and S. Ali, "An ethnobotanical survey of antidiabetic plants used by Hausa–Fulani tribes in Sokoto, Northwest Nigeria," *Journal of Ethnopharmacology*, vol. 172, pp. 91–99, 2015.
- [28] R. M. Al-Kindi, M. Al-Mushrafi, M. Al-Rabaani, and I. Al-Zakwani, "Complementary and alternative medicine use among adults with diabetes in Muscat region, Oman," *Sultan Qaboos University Medical Journal*, vol. 11, no. 1, pp. 62–68, 2011.
- [29] A. N. Kesari, S. Kesari, S. K. Singh, R. K. Gupta, and G. Watal, "Studies on the glycemic and lipidemic effect of *Murraya koenigii* in experimental animals," *Journal of Ethnopharmacology*, vol. 112, no. 2, pp. 305–311, 2007.
- [30] L. Dey, A. S. Attele, and C.-S. Yuan, "Alternative therapies for type 2 diabetes," *Alternative Medicine Review: A Journal of Clinical Therapeutic*, vol. 7, no. 1, pp. 45–58, 2002.
- [31] C. E. Momo, J. E. Oben, D. Tazoo, and E. Dongo, "Antidiabetic and hypolipidemic effects of *laportea ovalifolia* (urticaceae) in alloxan induced diabetic rats," *African Journal of Traditional, Complementary and Alternative Medicines*, vol. 3, no. 1, pp. 36–43, 2005.

- [32] C. W. Ssenyange, A. Namulindwa, B. Oyik, and J. Ssebuliba, "Plants used to manage type II diabetes mellitus in selected districts of central Uganda," *African Health Sciences*, vol. 15, no. 2, p. 496, 2015.
- [33] A. Meresa, W. Gemechu, H. Basha et al., "Herbal medicines for the management of diabetic mellitus in Ethiopia and Eretria including their phytochemical constituents," *AJADD*, vol. 5, no. 01, pp. 040–058, 2017.
- [34] A. Mohammed, C. D. Luka, A. L. Ngwen, O. F. R. Omale, and B. J. Yaknan, "Evaluation of the effect of aqueous leaf extract of jute mallow corchorus olitorius on some biochemical parameters in alloxan-induced diabetic rats," *European Journal of Pharmaceutical and Medical research*, vol. 6, no. 10, pp. 652–658, 2019.
- [35] K. Das, R. Tiwari, and D. Shrivastava, "Techniques for evaluation of medicinal plant products as antimicrobial agent: current methods and future trends," *Journal of Medicinal Plants Research*, vol. 4, no. 2, pp. 104–111, 2010.
- [36] J. Parekh and S. Chanda, "In vitro antimicrobial activity and phytochemical analysis of some Indian medicinal plants," *Turkish Journal of Biology*, vol. 31, no. 1, pp. 53–58, 2007.
- [37] E. Shellard, "The significance of research into medicinal plants," in *African Medicinal Plants, Proceeding of a Conference* University of Ife press, 1979.
- [38] A. Sofowora, *Screening Plants for Bioactive Agents. Medicinal Plants and Traditional Medicinal in Africa*, Spectrum Books Ltd, Sunshine House, Ibadan, Nigeria, 1993.
- [39] W. C. Evans, *Trease and Evans' Pharmacognosy E-Book*, Elsevier Health Sciences, 2009.
- [40] G. Trease and W. Evans, *Pharmacognosy*, Bailliere Tindall, London, UK, 1989.
- [41] J. B. Harborne, "Phenolic compounds," in *Phytochemical Methods* Springer, Berlin, Germany, 1973.
- [42] S. M. Salman, "Preliminary phytochemical, essential element analysis and antimicrobial activities of ethanolic extract of *Lotus corniculatus*," *International Journal of Biosciences*, vol. 7, no. 2, pp. 106–115, 2015.
- [43] P. Talele, "Isolation of starch from Ginger rhizome (*Zingiber officinale*)," *Journal of Pharmacognosy and Phytochemistry*, vol. 3, no. 6, pp. 157–162, 2015.
- [44] O. In, "Acute oral toxicity-Acute oral toxic class method. Guideline 423," *Eleventh Addendum to the OECD Guidelines for the Testing*, 2001.
- [45] A. Mushtaq, S. Akbar, M. A. Zargar et al., "Phytochemical screening, physicochemical properties, acute toxicity testing and screening of hypoglycaemic activity of extracts of *Eremurus himalaicus baker* in normoglycaemic Wistar strain albino rats," *BioMed Research International*, vol. 2014, pp. 1–6, 2014.
- [46] A. Sadat, M. Hore, K. Chakraborty, and S. Roy, "Phytochemical analysis and antioxidant activity of methanolic extract of leaves of corchorus olitorius," *International Journal of Current Pharmaceutical Research*, vol. 9, no. 5, pp. 59–63, 2017.
- [47] S. Parvin, M. Marzan's, S. Rahman, A. Kumer Das, S. Haque, and M. Rahmatullah, "Preliminary phytochemical screening, antihyperglycemic, analgesic and toxicity studies on methanolic extract of aerial parts of *Corchorus olitorius* L.," *Journal of Applied Pharmaceutical Science*, vol. 5, no. 9, pp. 68–71, 2015.
- [48] A. Adjatin, A. Hounkpatin, F. Assogba et al., "Phytochemical screening, antioxidant and cytotoxic activity of different morphotypes of *Corchorus olitorius* L. leaves in the central region of Benin Republic (West Africa)," *Journal of Pharmacognosy and Phytotherapy*, vol. 10, no. 12, pp. 195–203, 2018.
- [49] D. Orieko, O. Ohaeri, I. Ijeh, and S. Ijioma, "Identification of phytocomponents and acute toxicity evaluation of *Corchorus olitorius* leaf extract," *European Journal of Medicinal Plants*, vol. 23, pp. 1–16, 2018.
- [50] J. A. Saliu, G. Oboh, M. R. Schetinger, N. Stefanello, and J. B. T. Rocha, "Antidiabetic potentials of jute leaf (*Corchorus olitorius*) on type-2 diabetic rats," *Journal of Emerging Trends in Engineering and Applied Sciences*, vol. 6, no. 7, pp. 223–230, 2015.
- [51] S. H. Tiong, C. Looi, H. Hazni et al., "Antidiabetic and antioxidant properties of alkaloids from *Catharanthus roseus* (L.) G. Don," *Molecules*, vol. 18, no. 8, pp. 9770–9784, 2013.