

RESEARCH

Open Access



# Promptness in seeking treatment from Village Health workers for children under five years with malaria, diarrhoea and pneumonia in rural southwestern Uganda

Edgar Mulogo<sup>1\*</sup>, Stephen Baguma<sup>2</sup>, Moses Ntaro<sup>1</sup>, Shem Bwambale<sup>3</sup>, Michael Matte<sup>2</sup>, Andrew Wesuta<sup>2</sup>, David Ayebare<sup>1</sup>, Fred Bagenda<sup>1</sup> and Peter Kawungezi<sup>1</sup>

## Abstract

**Background** Village Health Workers (VHWs) in Uganda provide treatment for the childhood illness of malaria, pneumonia, and diarrhoea through the integrated community case management (iCCM) strategy. Under the strategy children under five years receive treatment for these illnesses within 24 h of onset of illness. This study examined promptness in seeking treatment from VHWs by children under five years with malaria, pneumonia, and diarrhoea in rural southwestern Uganda.

**Methods** In August 2022, a database containing information from the VHWs patient registers over a 5-year study period was reviewed (2014–2018). A total of 18,430 child records drawn from 8 villages of Bugoye sub-county, Kasese district were included in the study. Promptness was defined a caregiver seeking treatment for a child from a VHW within 24 h of onset of illness.

**Results** Sixty-four percent (64%) of the children included in the study sought treatment promptly. Children with fever had the highest likelihood of seeking prompt treatment (aOR = 1.93, 95% CI 1.80–2.06,  $p < 0.001$ ) as compared to those with diarrhoea (aOR = 1.43, 95% CI 1.32–1.52,  $p < 0.001$ ) and pneumonia (aOR = 1.33, 95% CI 1.24–1.42,  $p < 0.001$ ).

**Conclusion** The findings provide further evidence that VHWs play a critical role in the treatment of childhood illness in rural contexts. However, the proportion of children seeking prompt treatment remains below the target set at the inception of the iCCM strategy, in Uganda. There is a need to continually engage rural communities to promote modification of health-seeking behaviour, particularly for children with danger signs. Evidence to inform the design of services and behaviour change communication, can be provided through undertaking qualitative studies to understand the underlying reasons for decisions about care-seeking in rural settings. Co-design with communities in these settings may increase the acceptability of these services.

**Keywords** Promptness, Treatment, Childhood illness, Community health worker

\*Correspondence:

Edgar Mulogo

emulogo2000@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

In 2019 an estimated 5.2 million children under 5 years died mostly from preventable and treatable causes. These include; malaria, diarrhoea and pneumonia that can be prevented or treated with access to quality care by a trained health provider when needed, among other interventions [1]. The World Health Organization (WHO) estimates that a total of 33 million cases of malaria, diarrhoea and pneumonia go untreated every year. Early or timely access to medical care is one of the core approach in elimination of these illnesses in sub-Saharan Africa [2]. In order to decrease severity of childhood illnesses and its subsequent death, there is need for improving access to health workers [3] and appropriate health care-seeking behaviour of caregivers [4].

Diarrhoea, malaria, and pneumonia are the leading causes of mortality in children under-5 years of age in Africa [5, 6]. Sub-Saharan Africa and South Asia have registered the lowest reductions in mortality rates as more than 7 million children under age five (“U5s”) die every year of pneumonia, malaria and diarrhoea. In 2012, these regions contributed about 82% of the global under-five mortality. The two leading death causes of under-five children is pneumonia, accounting for 18% followed by diarrhoeal accounting for 15% of the total under-five mortality. For only malaria, over 90% globally occur in sub-Saharan Africa [7]. The region has age-long prevalence of under-5 mortalities despite increasing health intervention [8].

Evidence shows that most malaria related deaths in malaria affected countries occur at home without receiving appropriate or timely medical care, and when care is sought, it is often delayed [9]. Delay in seeking healthcare has been shown in several regions to play an important role in under-five morbidity and mortality [10]. A number of studies conducted in developing countries have also shown that delay in seeking appropriate care or not seeking any care causes a large number of child deaths [11]. Late care-seeking contributed to high mortality from acute respiratory infections in Uganda and from malaria in Tanzania. Even when drugs are free at community level caregivers do not seek treatment in time [12]. Other caregivers on contrary first use herbs and after this treatment fails they take their child to health workers when the situation has worsened [13]. In sub-Saharan Africa, responses to seek care outside home are often delayed, and when a response is made is at village drug shops characterized by inadequate trained workers [14, 15].

In Uganda alone, about 60% of parents with febrile children first seek care in the private sector [16, 17]. In most rural areas caregivers hardly recognize signs of

childhood illness for timely care-seeking [18]. Delay in seeking health care for a sick infant has been attributed to several factors for example combining home remedies with conventional treatments, inability to identify life-threatening illnesses and lack of knowledge. These challenges exist against a background of undiagnosed serious life threatening illness such as diarrhoea, and malaria [19].

In recent years, there has been a resurgence of interest by the international community in the use of VHWs as a way to bridge the gap in reducing child mortality through introducing iCCM as a way of timely handling of cases at village level. In iCCM model, trained VHWs are typically equipped to timely assess and treat the leading infectious causes of child mortality under 5 years [2, 20]. VHWs are trained, equipped with medicine and diagnostic gadgets to use at the community level. They also identify and refer children with severe disease to health facilities [21]. Government of Uganda with support from partners introduced iCCM to offer timely curative treatments for malaria, diarrhoea and pneumonia at community level. This was aimed at timely diagnosis and treatment to those who can hardly access health facilities [22]. However, the success of iCCM solidly depends on caregivers who should timely be able to recognize symptoms, seek health care from CHWs and adhere to treatment or referral within 24 h.

Despite progress made in reducing child mortality in Uganda, some caregivers’ choice for seeking and utilization of health care provided by community health workers is associated with severity of illness [23]. iCCM has the potential to increase access to prompt care, thereby decreasing morbidity and mortality [24]. A key objective of the iCCM programme in Uganda was to increase to at least 80% the proportion of children under-five years receiving appropriate treatment for malaria, pneumonia and diarrhoea within 24 h of onset of illness [25].

However, there is limited information on the promptness in seeking health treatment from VHWs, for children under five years with malaria, pneumonia and diarrhoea. This study, therefore, examined the promptness in seeking treatment for children under five years in a rural community.

## Methods

### Design and setting

A retrospective data review of records of VHWs in 8 villages of Bugoye sub-county was conducted in 2022. The review covered a period of five years (2014–2018). Bugoye Sub-county, Uganda, is a rural, mountainous area in Kasese District (on the western border of Uganda), VHWs from the national programme have provided

iCCM care since 2013, with financial and operational support from a long-standing collaboration with Mbarara University of Science and Technology (Mbarara, Uganda) and the Massachusetts General Hospital (Boston, Massachusetts, USA) [26].

The Bugoye VHWs were enlisted, trained and supplied to provide iCCM services in 22 villages of the sub-county, surround Bugoye Health Centre level III. Bugoye sub-county is home just over 45,000 people with an average household size of 5.6 people, with malaria and respiratory infections being the most predominant childhood illnesses. The VHW team is elected by its village community, with four to five VHWs derived from each of the villages. They receive comprehensive training to appropriately diagnose, treat and/or refer children under five years old with malaria, pneumonia or diarrhoea. Initially, VHWs participate in a basic 3-day training standardized by the Ugandan Ministry of Health (MoH) to review VHW responsibilities. This was followed with a 6-day training on implementing iCCM services [27].

The VHWs use the iCCM protocol, called the “Sick Child Job Aid”, to determine the proper care for each patient. They are equipped with rapid diagnostic tests (RDT) for malaria diagnosis in patients presenting with subjective fever; pneumonia is diagnosed based on age-based respiratory rate cut-offs, and diarrhoea is diagnosed by clinical history. For patients with “danger signs” (evidence of severe illness), VHWs provide initial assessment and referral or accompaniment to a health facility, as well as pre-referral treatment for some conditions. The five general danger signs include; vomiting everything, chest in-drawing, convulsions, not able to breast feed or drink, and very sleepy/unconscious/difficulty to wake [28]. VHWs use a “Sick Patient Register” to record each patient they assess. These Sick Patient Registers are submitted monthly and tallied to create a “Monthly Report” for the overall programme. The filed registers and monthly reports provided the data sources for this research.

#### Data collection

The filed registers provide the following information on the following; basic information about the patient (age, sex and presenting complaints), clinical assessment data (presence of danger signs, respiratory rate and malaria RDT result), actions taken (medications administered and/or referral to the health facility), as well as whether the patient was seen by the VHW within 24 h of onset of symptoms. An electronic database was created using de-identified data for all encounters from April 2014 to December 2018.

#### Variables and analysis

The study variables were derived from the information provided in the Sick Patient Registers. These variables include; sex, age, RDT result, whether the child had; diarrhoea, fast breathing, fever, danger sign, or was treated within 24 h. “Promptness” was defined as children seeking treatment from a VHW within 24 h of onset of illness (malaria, pneumonia, diarrhoea).

The data analysis was conducted using STATA version 14 software [29]. Two-sided chi-square tests for association were computed to detect differences between proportions of categorical variables such as sex, presenting compliant, presence of danger signs, referral, and whether patient visited the VHW within 24 h of onset of illness. The means of continuous variables such as age were compared using t-tests. A sliding scale of p-values was used as a basis to interpret the strength of association.

Logistic regression models were run to investigate the relationship between the outcome variable (promptness) and other variables. The model building strategy was not only limited to significant variables from the bivariate analysis, but also included independent variables that were considered to have social plausibility to promptness in seeking care [30].

## Results

### Background characteristics of the children

A total of 18,430 child encounters were analysed. Of these 11,724 (64%) visited the CHWs within 24 h. The background characteristics of the children are shown in Table 1. The majority of the children that had fever, diarrhoea and pneumonia visited the VHWs within 24 h of onset of illness ( $p < 0.05$ ). Only 37% of the children with danger signs visited the VHWs within 24 h of onset of illness ( $p < 0.001$ ).

### Factors associated with promptness in seeking care from VHWs

The factors associated with promptness in seeking care from VHWs are shown in Table 2.

## Discussion

The study findings show that the majority of children sought treatment from the VHWs within 24 h of illness. This is consistent with findings elsewhere that have reported promptness in seeking treatment from VHWs [28, 31]. Prompt seeking of treatment from the VHWs has been associated with short distance to reach the VHW, where the mother was the decision maker in the household, high trust and satisfaction with the services provided [28, 32]. The proportions of children with fever, diarrhoea and pneumonia who

**Table 1** Background characteristics of children

Characteristic	All (N = 18,430) n (%)	Visited VHW within 24 h		p-value
		No (6706) n (%)	Yes (11,724) n (%)	
Mean age in months (SD)	28.1 (± 17.1)	27.6 (± 17.6)	28.3 (± 16.9)	0.004 <sup>***</sup>
Sex (n = 18,297)				0.428
Female	9,105 (50)	3,266 (36)	5,839 (64)	
Male	9,192 (50)	3,349 (36)	5,843 (64)	
Fever (n = 18,430)				< 0.001 <sup>**</sup>
No	7,618 (41)	3,314 (44)	4,304 (56)	
Yes	10,812 (81)	3,392 (31)	7,420 (69)	
Diarrhoea (n = 18,430)				< 0.001 <sup>**</sup>
No	13,260 (72)	4,978 (38)	8,282 (62)	
Yes	5,170 (28)	1,728 (33)	7,420 (67)	
Pneumonia (n = 18,430)				0.026 <sup>*</sup>
No	10,091 (55)	3744 (37)	6347 (63)	
Yes	8339 (45)	2962 (36)	5377 (64)	
Child had danger sign (n = 18,430)				< 0.001 <sup>***</sup>
No	18,201 (99)	6561 (36)	11,640 (99)	
Yes	229 (1)	145 (63)	84 (37)	

SD standard deviation, † t test

\*significant at p < 0.05, \*\*significant at p < 0.01 and \*\*\*significant at p < 0.001

**Table 2** Association between patient characteristics and promptness in seeking care

Characteristics	Visit CHWs within 24 h		OR (95% CI)	p-value	aOR (95% CI)	p-value
	No	Yes				
Age						
< 12 months	1415	2060	1	< 0.001 <sup>***</sup>		
≥ 12 months	5291	9664	1.25 (1.16–1.35)		1.16 (1.07 – 1.25)	< 0.001 <sup>***</sup>
Sex						
Male	3349	5843	1	0.42		
Female	3266	5839	1.02 (0.96–1.08)		1.01 (0.95 – 1.07)	0.629
Fever						
No	3314	4304	1	< 0.001 <sup>***</sup>		
Yes	3392	7420	1.68 (1.58–1.78)		1.93 (1.80 – 2.06)	< 0.001 <sup>***</sup>
Diarrhoea						
No	4978	8282	1	< 0.001 <sup>***</sup>		
Yes	1728	3442	1.19 (1.11–1.28)		1.43 (1.32 – 1.54)	< 0.001 <sup>***</sup>
Pneumonia						
No	3744	6347	1	0.02 <sup>**</sup>		
Yes	2962	5377	1.07 (1.00–1.13)		1.33 (1.24 – 1.42)	< 0.001 <sup>***</sup>
Child had danger sign						
No	6561	11,640	1	< 0.001 <sup>***</sup>		
Yes	143	84	0.32 (0.24–0.42)		0.51 (0.38 – 0.69)	< 0.001 <sup>***</sup>
Child referred						
No	6124	11,212	1	< 0.001 <sup>***</sup>		
Yes	576	510	0.48 (0.42–0.54)		0.59 (0.52 – 0.68)	< 0.001 <sup>***</sup>

The patient characteristics independently associated with promptness in seeking care from a VHW are; fever (aOR = 1.93, 95% CI 1.80–2.06, p < 0.001), diarrhoea (aOR = 1.43, 95% CI 1.32–1.54, p < 0.001), pneumonia (aOR = 1.33, 95% CI 1.24–1.42, p < 0.001), child had danger sign (aOR = 0.51, 95% CI 0.38–0.69, p < 0.001), and child referred (aOR = 1.59, 95% CI 0.52–0.68, p < 0.001)

\*\*significant at p < 0.01 and \*\*\*significant at p < 0.001

sought care from the VHWs within 24 h of onset of symptoms was similar, but much lower than the national average at the time [25]. This may be attributed to the rugged mountainous terrain in which the communities dwell in comparison to other areas of the country, that may result in delays by some caregivers in seeking care for the children.

Children with fever sought treatment more promptly in comparison to those with diarrhoea and pneumonia. However studies elsewhere have reported that children with fast breathing seek treatment more promptly than those with either fever or diarrhoea [28, 31]. This study noted that children with danger signs were less likely to seek treatment promptly. It is assumed that children with danger signs, which often reflects more severe illness would be expected to seek treatment promptly. However, it has been reported that caregivers often do not recognize symptoms and danger signs of severely ill children, which may lead to delays in seeking treatment [33, 34]. On the other hand, a number of children with danger signs bypass the VHWs and go directly to the health facilities which is the appropriate action.

### Limitations

Although the study data is from an iCCM programme in one sub county the context and the providers are typical of other rural setting in Uganda. The study findings are therefore comparable across similar settings in the country. The study data is also cross-sectional in nature therefore a temporal relationship between the independent variables and outcome cannot be elucidated. The direction of causality can therefore only be regarded as suggestive.

### Conclusions

The findings provide further evidence that VHWs play a critical role in the treatment of childhood illness in rural contexts. However, the proportion of children seeking prompt treatment remains below the target set at the inception of the iCCM strategy, in Uganda. There is a need to continually engage rural communities to promote modification of health-seeking behaviour, particularly for children with danger signs. Evidence to inform the design of services and behaviour change communication, can be provided through undertaking qualitative studies to understand the underlying reasons for decisions about care-seeking in rural settings. Co-design with communities in these settings may increase the acceptability of these services.

### Acknowledgements

The study team acknowledges the support and contributions of the; Massachusetts General Hospital, Department of Medicine MGH, Bugoye

Health Centre Level III, VHWs in Bugoye Sub-county, Moses Wetyanga, Sarah Masika, Bwambale Aprunale and Mbusa Rapheal.

### Author contributions

SB, MN, SB, MM, AW, DA, FB and PK were involved in the conception and design of the study, and its implementation. EM participated in the conception and design of the study, analysis of the data, interpretation of the findings and drafting of the paper.

### Funding

This work was supported by the Mooney-Reed Charitable Foundation and the Massachusetts General Hospital Department of Medicine.

### Availability of data and materials

All data supporting our findings are contained in the paper. There are no restrictions to data sources, however, details of the full data may be accessed through Prof. Edgar Mugema Mulogo (corresponding author), Department of Community Health, Mbarara University of Science and Technology, PO Box 1410, Mbarara, Uganda.

### Declarations

#### Ethics approval and consent to participate

Ethical approval for this study was sought and obtained from the Research Ethics Committee at Mbarara University of Science and Technology. Written informed consent was also obtained from the individual subjects.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that there are no competing interests.

#### Author details

<sup>1</sup>Department of Community Health, Mbarara University of Science and Technology, P.O. Box 1410, Mbarara, Uganda. <sup>2</sup>Bugoye Community Health Collaboration, P.O. Box 149, Kasese, Uganda. <sup>3</sup>Bugoye Health Centre, Kasese District Local Government, P.O. Box 149, Kasese, Uganda.

Received: 8 September 2022 Accepted: 21 June 2023

Published online: 27 June 2023

### References

1. WHO. Children: improving survival and well-being. Geneva: World Health Organization; 2020. <https://www.who.int/news-room/fact-sheets/detail/children-reducing-mortality>. Accessed 8 Sept 2020.
2. Young M, Wolfheim C, Marsh DR, Hammamy D. World Health Organization/United Nations Children's Fund joint statement on integrated community case management: an equity-focused strategy to improve access to essential treatment services for children. *Am J Trop Med Hyg*. 2012;87(Suppl 5):6–10.
3. WHO. Global strategy for women's, children's and adolescents' health. Geneva: World Health Organization; 2016.
4. Adedire EB, Asekun-Olarinmoye EO, Fawole O. Maternal perception and care-seeking patterns for childhood febrile illnesses in rural communities of Osun state, South-Western Nigeria. *Sci J Public Health*. 2015;2:636–43.
5. Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, Bassani DG, et al. Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet*. 2010;375:1969–87.
6. Liu L, Johnson HL, Cousens S, Perin JSS, Lawn JE, et al. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet*. 2012;379:2151–61.
7. WHO, UNICEF. UNICEF joint statement: management of pneumonia in community settings. Geneva: World Health Organization; 2014.
8. Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J, et al. Global, regional, and national causes of under-5 mortalities in 2000–15: an updated systematic analysis with implications for the Sustainable Development Goals. *Lancet*. 2016;388:3027–35.



9. Wiseman V, Scott A, Conteh L, McElroy B, Stevens W. Determinants of provider choice for malaria treatment: experiences from The Gambia. *Soc Sci Med*. 2008;67:487–96.
10. Rutebemberwa E, Buregyeya E, Lal S, Clarke SE, Hansen KS, Magnussen P, et al. Assessing the potential of rural and urban private facilities in implementing child health interventions in Mukono district, central Uganda—a cross sectional study. *BMC Health Serv Res*. 2016;16:268.
11. Amarasiri de Silva M, Wijekoon A, Hornik R, Martines J. Care seeking in Sri Lanka: one possible explanation for low childhood mortality. *Soc Sci Med*. 2001;53:1363–72.
12. Nsungwa-Sabiiti J, Peterson S, Pariyo G, Ogwal-Okeng J, Petzold MG, Tomson G. Home-based management of fever and malaria treatment practices in Uganda. *T Trans R Soc Trop Med Hyg*. 2007;101:199–207.
13. Diaz T, George AS, Rao SR, Bangura PS, Baimba JB, McMahan SA, Kabano A. Health care seeking for diarrhea, malaria and pneumonia among children in four poor rural districts in Sierra Leone in the context of free health care results of a cross-sectional survey. *BMC Public Health*. 2013;13:157.
14. Kolola T, Gezahegn T, Addisie M. Health care seeking behavior for common childhood illnesses in Jeldu District, Oromia Regional State, Ethiopia. *PLoS ONE*. 2016;11: e0164534.
15. Mbonye AK, Buregyeya E, Rutebemberwa E, Clarke SE, Lal S, Hansen KS, et al. Referral of children seeking care at private health facilities in Uganda. *Malar J*. 2017;16:76.
16. Rutebemberwa E, Pariyo G, Peterson S, Tomson G, Kallander K. Utilization of public or private health care providers by febrile children after user fee removal in Uganda. *Malar J*. 2009;8:45.
17. Rutebemberwa E, Kallander K, Tomson G, Peterson S, Pariyo G. Determinants of delay in care-seeking for febrile children in eastern Uganda. *Trop Med Int Health*. 2009;14:472–9.
18. Pérez-Cuevas R, Guiscafré H, Romero G, Rodríguez L, Gutierrez G. Mother's health seeking behavior in acute diarrhea in Tlaxcala, Mexico. *J Diarrhoeal Dis Res*. 1996;14:260–8.
19. WHO. Integrated management of childhood illnesses. Module 5. Geneva: World Health Organization; 2015. [http://www.who.int/maternal\\_child\\_adolescent/topics/child/imci/en/](http://www.who.int/maternal_child_adolescent/topics/child/imci/en/). Accessed 22 Feb 2021.
20. WHO. Towards a grand convergence for child survival and health: a strategic review of options for the future building on lessons learnt from IMNCI. Geneva, World Health Organization. 2016. <http://www.who.int/maternal-child-adolescent/documents/strategic-review-child-health-imnci/en/>. Accessed 22 Feb 2021.
21. Rowe AK, Rowe SY, Snow RW, Korenromp EL, Armstrong-Schellenberg JR, Stein C, et al. The burden of malaria mortality among African children in the year 2000. *Int J Epidemiol*. 2006;35:691–704.
22. Winch PJ, Gilroy KE, Wolfheim C, Starbuck ES, Young MW, Walker LD, et al. Intervention models for the management of children with signs of pneumonia or malaria by community health workers. *Health Policy Plan*. 2005;20:199–212.
23. WHO. Causes of child mortality 2016. Geneva: World Health Organization; 2016.
24. Miller JS, English L, Matte M, Mbusa R, Ntaro M, Bwambale S, et al. Quality of care in integrated community case management services in Bugoye, Uganda: a retrospective observational study. *Malar J*. 2018;17:99.
25. Ministry of Health. Integrated community case management of childhood malaria, pneumonia and diarrhoea: implementation guidelines. Kampala: MOH/WHO/UNICEF; 2010.
26. Miller JS, Mulogo EM, Wesuta AC, Mumbere N, Mbaju J, Matte M, et al. Long-term quality of integrated community case management care for children in Bugoye Subcounty, Uganda: a retrospective observational study. *BMJ Open*. 2022;12: e051015.
27. English L, Miller JS, Mbusa R, Matte M, Kenney J, Bwambale S, et al. Monitoring iCCM referral systems: Bugoye Integrated Community Case Management Initiative (BIMI) in Uganda. *Malar J*. 2016;15:247.
28. Ministry of Health. ICCM Facilitator Guide: Caring for Newborns and Children in the Community. Kampala: Ministry of Health; 2010.
29. StataCorp L. StataCorp: stata statistical software: release 14. College Station: StataCorp LP; 2015.
30. Victora CG, Huttly SR, Fuchs SC, Olinto MT. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. *Int J Epidemiol*. 1997;26:224–7.
31. Kalyango JN, Alfven T, Peterson S, Mugenyi K, Karamagi C, Rutebemberwa E. Integrated community case management of malaria and pneumonia increases prompt and appropriate treatment for pneumonia symptoms in children under five years in Eastern Uganda. *Malar J*. 2012;12:340.
32. Mazzi M, Bajunirwe F, Aheebwea E, Nuwamanya S, Bagenda FN. Proximity to a community health worker is associated with utilization of malaria treatment services in the community among under-five children: a cross-sectional study in rural Uganda. *Int Health*. 2018;11:143–9.
33. Tuhebwe D, Tumushabe E, Leontsini E, Wanyenze RK. Pneumonia among children under five in Uganda: symptom recognition and actions taken by caretakers. *Afr Health Sci*. 2014;14:993–1000.
34. Ndu IK, Ekwochi U, Osuorah CD, Onah KS, Obuoha E, Odetunde OI, et al. Danger signs of childhood pneumonia: caregiver awareness and care seeking behavior in a developing country. *Int J Pediatr*. 2015;2015: 167261.

### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Ready to submit your research? Choose BMC and benefit from:**

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

**At BMC, research is always in progress.**

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)



## Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

[onlineservice@springernature.com](mailto:onlineservice@springernature.com)