



# HIV, Social Networks, and Loneliness among Older Adults in Uganda

Lien T. Quach<sup>1,2,3,4,5,16</sup> · Christine S. Ritchie<sup>2,3,5</sup> · Zahra Reynolds<sup>1</sup> · Robert Paul<sup>6</sup> · Janet Seeley<sup>7</sup> · Yao Tong<sup>1</sup> · Susanne Hoepfner<sup>5,8</sup> · Samson Okello<sup>9,10</sup> · Noeline Nakasujja<sup>11</sup> · Brianne Olivieri-Mui<sup>12</sup> · Deanna Saylor<sup>13</sup> · Meredith Greene<sup>14</sup> · Stephen Asiimwe<sup>1,9,15</sup> · Edna Tindimwebwa<sup>15</sup> · Flavia Atwiine<sup>9</sup> · Ruth Sentongo<sup>9</sup> · Mark J. Siedner<sup>1,5,9</sup> · Alexander C. Tsai<sup>5,8,9</sup>

Accepted: 26 December 2023

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2024

## Abstract

Loneliness among older adults has been identified as a major public health problem. Yet little is known about loneliness, or the potential role of social networks in explaining loneliness, among older people with HIV (PWH) in sub-Saharan Africa, where 70% of PWH reside. To explore this issue, we analyzed data from 599 participants enrolled in the Quality of Life and Ageing with HIV in Rural Uganda study, including older adults with HIV in ambulatory care and a comparator group of people without HIV of similar age and gender. The 3-item UCLA Loneliness Scale was used to measure loneliness, and HIV status was the primary explanatory variable. The study found no statistically significant correlation between loneliness and HIV status. However, individuals with HIV had smaller households, less physical and financial support, and were less socially integrated compared to those without HIV. In multivariable logistic regressions, loneliness was more likely among individuals who lived alone (aOR:3.38, 95% CI:1.47–7.76) and less likely among those who were married (aOR:0.34, 95% CI:0.22–0.53) and had a higher level of social integration (aOR:0.86, 95% CI: 0.79–0.92). Despite having smaller social networks and less support, older adults with HIV had similar levels of loneliness as those without HIV, which may be attributed to resiliency and access to HIV-related health services among individuals with HIV. Nonetheless, further research is necessary to better understand the mechanisms involved.

**Keywords** HIV · Social Networks · Social Integration · Loneliness · Older Adults · Uganda

## Introduction

The average age of the population of people with HIV (PWH) is increasing, especially in sub-Saharan Africa. In 2020, approximately 70% of the 37 million people with HIV (PWH) worldwide lived in sub-Saharan Africa, and more than 50% were over 50 years of age [1]. Loneliness is prevalent among older people, particularly among older PWH in high-income settings [2], and is associated with poor control of chronic medical conditions, depression, and adverse health outcomes, including death. The magnitude of these associations is greater among older adults [3–5]. The experience of loneliness has been conceptualized as the feeling of distress that accompanies a perceived unmet need

for relationships, either in quantity or in quality [6]. Much of the literature on loneliness and health among older people has been derived from high-income countries, where feelings of loneliness are endorsed by approximately one-third of older adults [7], and nearly half of older PWH [2, 8]. Much less is known about loneliness in older PWH in sub-Saharan Africa.

The scarcity of data about the prevalence of loneliness among older people in sub-Saharan Africa highlights a significant research gap. Given that loneliness is a globally significant public health issue for older adults, it is essential to delve into this topic within specific contexts. The surge in urbanization, particularly in sub-Saharan Africa [9], coupled with changing family structures [10–12], may compromise social support systems, suggesting that loneliness might be a pressing concern for older individuals in this region [11, 12]. On the other hand, the persistence of multigenerational families [10], which have traditionally served as sources of intergenerational support [10], could

Mark J. Siedner and Alexander C. Tsai contributed equally to this work.

Extended author information available on the last page of the article

potentially alleviate loneliness. Furthermore, collectivist cultures characterized by values such as obligation, cohesion, and commitment might serve a protective function against loneliness [13]. Studies of older PWH in rural South Africa and Uganda have also shown that having a stronger, more extensive, and primarily family-based social network is associated with greater interaction with social network members and greater community interdependence [14]. Other studies have shown that PWH tend to exhibit resilience as a consequence of managing multiple chronic health conditions [15–17] and achieving mastery [18, 19]. Studies from South Africa have shown that older PWH have access to healthcare support (e.g., through clinics funded by the U.S. President’s Emergency Plan for AIDS Relief) that may not be available to HIV-negative older people [20, 21]. And finally, older PWH contend with unique challenges, including stigma, discrimination, smaller social networks, and social isolation [22]. These patterns underscore the need to comprehensively understand the factors that contribute to loneliness among older PWH.

The present study is motivated by Social Provisions Theory [23–30]. Through an examination of factors such as social networks, social integration, various types and sources of social support, and their differentiation between older adults living with and without HIV, this framework can help to understand the determinants of loneliness that can help trace their origins to family, friends, mentors/providers, or one’s own sense of self [23–27]. We hypothesize that these constructs are crucial for flourishing among older adults with chronic health conditions, including HIV, because they can help older adults adhere to treatment, maintain healthy lifestyles, cope with daily challenges, and alleviate loneliness [19, 31].

To address the gaps in the literature reviewed above, we conducted a cross-sectional analysis of baseline data from older people enrolled in an ongoing cohort study in rural Uganda. The primary objective of the present analysis was to estimate the association between HIV serostatus and loneliness. We also sought to compare the two groups in terms of social networks, social support, and social integration, in order to investigate how they affect loneliness in this setting.

## Methods

### Study Population and Data Collection

We analyzed data from 599 participants enrolled in the *Quality of Life and Ageing with HIV in Rural Uganda Study*, a study of quality of life among older PWH in a rural region of southwestern Uganda [32]. PWH were eligible to

participate if they were older than 49 years of age and had been on ART for at least three years and were engaged in ambulatory HIV care at the Mbarara Regional HIV Clinic or the Kabwohe Clinical Research Center HIV Clinic ( $n = 297$ ). We then recruited an age- and sex-similar group of people without HIV (PWOH) from census data located in the same clinic catchment areas [33]. Data were collected during October 2020–October 2021. Due to the COVID-19 epidemic, data were collected via phone interview during this wave of the study.

### Measures

Our primary explanatory variable of interest was HIV serostatus, which was based on confirmatory HIV testing [34]. The primary outcome was the 3-item UCLA Loneliness Scale [35], which queries participants about whether they “never,” “sometimes,” or “often” feel a lack of companionship, feel left out of community meetings or events, or feel isolated from others. We assigned one point to “never” responses, two points to “sometimes,” and three points to “often” responses, allowing for a total loneliness score ranging from 3 to 9, with higher scores indicating a higher degree of loneliness. We followed Steptoe et al. [36] in defining loneliness at the fifth quintile of the distribution of the total loneliness score ( $\geq 5$  in this study sample). The UCLA Loneliness Scale has been used in many African settings, including Uganda, South Africa and Zimbabwe and Ghana, and has shown consistent and reliable measurement properties across many countries [37].

To better understand similarities or differences in loneliness between PWH vs. PWOH, we compared the two groups on several measures of social connection factors based on Social Provisions Theory [25, 28–30], including social networks, social support and social integration:

- We measured three structural aspects of social networks by eliciting the number of people living in the respondent’s household, marital status (married or cohabitating with a partner vs. divorced/separated, widowed, or single), and whether the study participant reported living alone.
- We measured physical social support [38] by characterizing the *types* of physical support they received from others (e.g., buying food, agricultural work, fetching water, cooking, going to the clinic or traditional healer, and collecting firewood; maximum of 6 types) and the *sources* of familial physical support (spouse, parent, son/daughter, grandson, granddaughter, son/daughter-in-law, and other relatives; maximum of 4 sources). We also elicited whether they received physical support

from any community source (community members, neighbors, church attendees, or paid helpers).

- Similar to the above, we measured financial social support [38] by characterizing the *types* of financial support (e.g., paying for medicines, doctor visits, clinical or hospital fees, food, clothing, transportation, school expenses for offspring; maximum of 6 types), the *sources* of familial financial support (maximum of 3 sources as above), and receipt of any financial support from any community source.
- We measured social integration by assessing the total number of social groups in which each respondent participated. We took a comprehensive census of all community groups in the area and categorized them as follows [39]: vocational groups; positive living groups for PWH; local council committees; water committees; village health teams; National Agriculture Advisory Services groups; church or other religious groups; women's groups; gardening committees; community burial groups; clan groups; and revolving funds, savings and credit co-operative society (SACCO) groups, registered savings groups, or other community or village savings groups. The total social integration score was the total number of groups in which the respondent reported participation (out of a maximum of 20).

Additional covariates included age; sex; educational attainment; self-reported alcohol consumption (never vs. any); and the number of self-reported comorbidities, including diabetes, high blood pressure, heart attack or heart failure, kidney problems, stroke, cancer, chronic obstructive pulmonary diseases, asthma, pneumonia, high cholesterol, and tuberculosis.

## Statistical Analysis

We compared PWH vs. PWOH on the loneliness and social connection variables, using Student's *t*-tests, log-rank tests, and chi-squared tests as appropriate. To estimate the association between HIV status and loneliness, we fitted a multivariable logistic regression model with loneliness specified as the outcome and HIV as the primary explanatory variable, while adjusting for the covariates listed above. As described in more detail below, we unexpectedly observed statistically significant differences between PWH and PWOH on several of the social connection variables but no statistically significant difference on the primary outcome. We therefore estimated a series of multivariable logistic regression models specifying loneliness as the outcome and the social connection variables as the explanatory variables, with each regression model including one of the social connection variables and adjusting for the covariates listed above (10 regression

models in total). To assess the robustness of our findings to misclassification resulting from the potentially arbitrary cut-off threshold in the outcome, we also fit a series of negative binomial regressions specifying the total loneliness score (ranging from 3 to 9) as a count outcome variable. Statistical significance was designated at the conventional 0.05 level. All analyses were conducted with SAS version 9.4 [40].

## Ethical Considerations

The study was approved by the ethics committees at Mass General Brigham in the United States and at the Mbarara University of Science and Technology in Uganda. We also obtained clearance to conduct the study from the Uganda National Council of Science and Technology. All participants consented to participate in the study verbally. The review committees waived written consent due to the COVID-19 pandemic and the infeasibility of obtaining written consent during remote data collection.

## Results

The study sample included 297 PWH and 302 PWOH (Table 1). The mean age of participants was 58 years (range, 49–88 years). By design, women accounted for 49% of the study population (295/599). Most study participants had achieved a primary school level of education or less (74%, 442/599). The mean number of comorbidities was 0.5 (standard deviation [SD] 0.8; range, 0–5). PWH were less likely to be married or report alcohol use, but otherwise there were no statistically significant differences in age, sex, education, or comorbidities between PWH and PWOH.

PWH had smaller household sizes (3.5 [SD, 2.2] vs. 3.9 [SD, 2.1],  $Z=-2.7$ ,  $P=0.01$ ) and were more likely to live alone (7.1% [21/297] vs. 1.3% [4/302], Fisher's exact test  $P<0.001$ ). PWH had lower social integration scores (2.8 [SD, 3.1] vs. 3.8 [SD, 3.3],  $Z=-4.1$ ,  $P<0.001$ ) (Fig. 1A), received fewer types of physical support (2.3 [SD, 1.9] vs. 3.3 [SD, 1.8],  $Z=-6.4$ ,  $P<0.001$ ) (Fig. 1B) and fewer sources of physical support (1.1 [SD, 1.2] vs. 1.6 [SD, 1.1],  $Z=-2.8$ ,  $P<0.001$ ) (Fig. 1C). They also had fewer types of financial support (0.9 [SD, 1.8] vs. 1.3 [SD, 2.0],  $Z=-5.0$ ,  $P<0.001$ ) (Fig. 1D), and fewer sources of financial support (0.3 [SD, 0.6] vs. 0.4 [SD, 0.6],  $Z=-2.9$ ,  $P<0.001$ ) (Fig. 1E). We found no statistically significant differences in the receipt of community physical support and community financial support.

A substantial proportion of study participants reported feeling lonely: 179 (29.8%) felt they "sometimes"/"often" lacked companionship, 158 (26.3%) felt left out of community meetings, and 118 (19.7%) felt isolated from others.


**Table 1** Characteristics of the sample, stratified by HIV Status

Characteristics	Total (N= 599)	HIV-negative (N=302)	HIV-positive (N=297)	Statistic's Values	Z, chi-square, or Fisher
Age (years), mean [standard deviation (SD)]	58.4 (6.6)	58.47 (6.8)	58.34(6.4)	-0.02	$\alpha$
Women, N (%)	295 (49.2)	148 (49.0)	147 (49.5)	0.01	$\infty$
Educational attainment, N (%)					
Primary or less (no school or P1-P7)	442 (73.8)	226 (74.8)	216 (72.7)	0.45	$\infty$
Secondary (S1-S6)	95 (15.9)	47 (15.6)	48 (16.2)		
Post-secondary	62 (10.4)	29 (9.6)	33 (11.1)		
Alcohol consumption, N (%)	143 (23.7)	82 (27.2)	60 (20.2)	3.40	$\infty$
Total comorbidities, mean (SD)	0.5 (0.8)	0.5 (0.8)	0.5 (0.8)	0.13	$\alpha$
Household composition					
Son, N (%)	287 (47.9)	151 (50.0)	136 (45.8)	1.06	$\infty$
Daughter, N (%)	279 (46.6)	149 (49.3)	130 (43.8)	1.86	$\infty$
Non-biological dependent child, N (%)	14 (2.3)	6 (2)	8 (2.7)	0.33	$\infty$
Grandson, N (%)	184 (30.7)	97 (32.1)	87 (29.3)	0.56	$\infty$
Granddaughter, N (%)	162 (27)	88 (29.1)	74 (24.9)	1.35	$\infty$
Paid helper, N (%)	47 (7.8)	21 (7.0)	26 (8.8)	0.67	$\infty$
Parent, N (%)	24 (4)	13 (4.3)	11 (3.7)	0.14	$\infty$
Nephew, N (%)	17 (2.8)	7 (2.3)	10 (3.4)	0.60	$\infty$
Niece, N (%)	13 (2.2)	7 (2.3)	6 (2.0)	0.06	$\infty$
Other family members or relatives <sup>‡</sup> , N (%)	51(8.5)	22 (7.3)	29 (9.8)	1.18	$\infty$
Social networks					
Living with spouse/partner, N (%)	404 (67.4)	242 (80.1)	162 (54.5)	44.65 ***	$\infty$
Household size, mean (SD)	3.7 (2.2)	3.91 (2.1)	3.46 (2.2)	-2.70 **	$\alpha$
Living alone	25 (4.2)	4 (1.3)	21 (7.1)	0.0004 ***	$\Omega$
Types of physical support, mean (SD)	2.8 (1.9)	3.3(1.8)	2.3 (1.9)	-6.39 ***	$\alpha$
Number of familial sources of physical support, mean (SD)	1.3 (1.1)	1.6 (1.1)	1.1 (1.2)	-2.80 ***	$\alpha$
Any community source of physical support, N (%)	81 (13.5)	36 (11.9)	45 (15.2)	1.33	$\infty$
Types of financial support, mean (SD)	1.1 (1.9)	1.3 (2.0)	0.9 (1.8)	-5.01 **	$\infty$
Number of familial sources of financial support, mean (SD)	0.3 (0.6)	0.4 (0.6)	0.3 (0.6)	-2.93 **	$\infty$
Any community source of financial support, N (%)	8.0 (1.3)	4.0 (1.3)	4.0 (1.4)	0.27	$\Omega$
Social integration score, mean (SD)	3.3 (3.2)	3.77 (3.3)	2.8 (3.1)	-4.09 ***	$\infty$
Total loneliness score, mean (SD)	3.9 (1.4)	3.83 (1.3)	4.03 (1.4)	2.15 *	$\infty$
Classified as "lonely", N (%)	156 (26.0)	72 (23.8)	84 (28.3)	5.40	$\infty$

**Notes:** <sup>‡</sup>Other family members and other relatives such as brother, sister, sister-in-law, parent-in-law, son-in-law, brother-in-law, daughter-in-law, and other relatives

<sup>‡</sup>Z score of the Wilcoxon Two-Sample Test for continuous or count variables; <sup>‡</sup>Chi-square test for binary or categorical variables; <sup>‡</sup>Fisher's exact test

\* $p \leq 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ;



**Fig. 1** Comparison of social integration, physical, and financial support among people with HIV (PWH) vs. people without HIV

Altogether 156 (26.0%) met the threshold definition of loneliness, with a score of 5 or greater on the loneliness scale. A higher proportion of PWH (vs. PWOH) met the threshold definition of loneliness, but in contradiction of our hypothesis based on existing literature, the comparison with PWOH did not reveal statistically significant differences in loneliness by HIV serostatus (28.3% vs. 23.8%, chi-square = 5.4,  $P=0.22$ ). This finding persisted after multivariable adjustment (adjusted odds ratio [aOR]: 1.24, 95% CI: 0.86–1.80, Table 2). In a series of multivariable logistic regression models in which we specified the social connection variables as the primary explanatory variables of interest while adjusting for sociodemographic covariates (10 regression models in all), loneliness was inversely associated with living with a spouse/partner (aOR: 0.34, 95% CI: 0.22–0.53) and with greater social integration (aOR: 0.86 per community group, 95% CI: 0.81–0.91), but positively associated with living alone (aOR: 3.38, 95% CI: 1.47–1.86). Loneliness did not have a statistically significant association with any of the physical or financial support variables (Table 2).

The results from negative binomial regressions, in which the total loneliness score was specified as a count variable, indicated similar findings for the association between HIV and loneliness, and between several of the social connection variables and loneliness (Table 3).

## Discussion

In a cross-sectional study of 599 older PWH in rural Uganda, and an age- and sex-similar sample of PWOH, we found no statistically significant difference in loneliness between

the two groups. This finding differs from similar studies in high-income settings, where PWH generally report a higher prevalence of loneliness than PWOH [2]. Our finding was unexpected, particularly given that we did find that PWH generally had a restricted range of social connections: they were more likely to live alone, were less likely to live with a spouse or primary partner, had smaller household sizes, reported fewer types and sources of support, and were less socially integrated. In this regard, our findings are consistent with prior work [41]. Considered in light of other recent findings from this same cohort showing that PWH had higher health-related quality of life compared with PWOH [32, 42], these findings suggest a certain degree of resilience among PWH despite structural disadvantages in the nature and breadth of their social connections [43–46]. This assumption is supported by a previous study's finding of multiple resilience resources among PWH that helps them manage their health conditions and improve their well-being in order to survive and flourish [43–45, 47]. Furthermore, social support accessed through clinic-based medical care, such as antiretroviral therapy programs and peer support groups, could also play a role in enhanced resilience and better coping among PWH, helping them to reduce the impacts of stigma and discrimination and, thus, reduce loneliness among older PWH [43–45, 48]. Consistent with this idea, other studies of PWH in rural Uganda have shown that both depression and internalized stigma decline over time on antiretroviral therapy [37, 49, 50]. Future research should further investigate the impact of formal health services on various forms of resilience and explore the differences in traditional social cohesion across generations [51] and the effect of reduced cohabitation on access to and receipt of

**Table 2** Correlates of loneliness, specified as a binary dependent variable

Exposures	Unadjusted <sup>∇</sup>	Adjusted
	OR (95% CI)	A OR (95% CI)
HIV-positive	1.26 (0.87, 1.82)	1.27 (0.88, 1.86)
Lives alone	3.27 (1.46, 7.32)	3.38 (1.47, 7.76)
Married	0.29 (0.2, 0.42)	0.34 (0.22, 0.53)
Household size	0.95 (0.88, 1.04)	0.96 (0.88, 1.05)
Types of physical support	0.97 (0.88, 1.07)	1.003 (0.91, 1.11)
Familial sources of physical support	0.87 (0.74, 1.03)	0.92 (0.77, 1.09)
Any community source of physical support	0.66 (0.37, 1.19)	0.61 (0.33, 1.12)
Types of financial support	1.04 (0.95, 1.15)	0.98 (0.89, 1.08)
Familial sources of financial support	1.12 (0.83, 1.52)	0.92 (0.67, 1.27)
Any community source of financial support	1.72 (0.41, 7.28)	1.26 (0.29, 5.51)
Social integration	0.86 (0.8, 0.93)	0.85 (0.79, 0.92)

**Notes:** <sup>∇</sup>Each cell represents the output of a single regression model with loneliness specified as the outcome and the row variable specified as the primary explanatory variable of interest. Thus, the estimates in column 1 are derived from 11 univariable logistic regression models. The estimates in column 3 are derived from 11 multivariable logistic regressions that also include covariate adjustment for age, sex, education, alcohol consumption, and comorbidities

OR: Odds Ratio

CI: Confidence Interval

**Table 3** Correlates of loneliness, specified as a count variable

Exposures	Unadjusted <sup>∇</sup>	Adjusted
	IRR (95% CI)	IRR (95% CI)
HIV-positive	1.05 (0.97, 1.14)	1.05 (0.97, 1.14)
Lives alone	1.22 (1.02, 1.47)	1.21 (1.01, 1.46)
Married	0.80 (0.74, 0.87)	0.82 (0.74, 0.91)
Household size	0.99 (0.97, 1.00)	0.99 (0.97, 1.01)
Types of physical support	0.99 (0.97, 1.02)	0.9995 (0.98, 1.02)
Familial sources of physical support	0.98 (0.94, 1.01)	0.98 (0.95, 1.02)
Any community source of physical support	0.95 (0.84, 1.07)	0.93 (0.83, 1.06)
Types of financial support	1.00 (0.98, 1.03)	0.99 (0.97, 1.01)
Familial sources of financial support	1.00 (0.94, 1.08)	0.97 (0.90, 1.04)
Any community source of financial support	1.05 (0.74, 1.48)	0.997 (0.71, 1.41)
Social integration	0.98 (0.96, 0.99)	0.97 (0.96, 0.99)

**Notes:** <sup>∇</sup> Each cell represents the output of a single regression model with loneliness specified as the outcome and the row variable specified as the primary explanatory variable of interest. Thus, the estimates in column 1 are derived from 11 univariable negative binomial regression models. The estimates in column 3 are derived from a single multivariable negative binomial regression model that also includes covariate adjustment for age, sex, education, alcohol consumption, and comorbidities

IRR: Incidence Rate Ratio

CI: Confidence Interval

care and support, particularly as PWH grow into older age [52].

Notwithstanding the relatively smaller social networks among PWH, most study participants in our study in Uganda lived with extended family, and less than 10% lived alone. This finding is consistent with previous studies conducted in Uganda and other countries throughout sub-Saharan Africa, which have demonstrated how family structures can serve as a *de facto* social security system and compensate for the often limited formal services available for aging populations [47, 51, 53]. These social arrangements contrast starkly with data from resource-rich settings. However, a previous household survey in Uganda found that older people living alone lacked family support networks and formal community care centers [51]. Because traditional social cohesion is changing across generations as a result of immigration and economic development, there is a need to strengthen the social support system for older adults in their later years, especially older PWH living alone [51, 54]. The more expansive social networks, stronger kinship ties, and greater community interdependence in rural areas of Africa likely have an impact on the needs of PWH as they age, with concomitant implications for the support services needed for this population.

Two additional findings from our study are worthy of note. First, we found that greater social integration is inversely associated with loneliness. This finding, which is consistent with reporting from high-income countries [55], may strengthen the case for the benefit of involvement in social activities, which increases social capital, positive health outcomes, and overall well-being in older adults

in the region as well [56]. In addition, a previous study showed that social integration is associated with a sense of belonging because it allows older adults to engage with and expand their social networks and to feel more loved [57]. Interventions to strengthen the social integration of older-age people in rural Uganda may reduce loneliness among both PWH and PWOH. Second, we found that loneliness was more prevalent among women. Prior studies on sex differences in loneliness have yielded mixed findings. Two studies of older adults in the global north similarly showed that loneliness was more prevalent among women [58–60], potentially due to longer life expectancies and therefore greater risks for widowhood, living alone, chronic illness, disability, and functional limitations, all of which are likely associated with higher risk of depression in women in high income settings [59, 61]. In contrast, a study of community-dwelling older-age Mozambican migrants in South Africa found that loneliness was more prevalent among men, due to their lower rates of social support, social participation and smaller social networks [62]. Future studies may identify specific psychosocial mechanisms to explain these gender differences in loneliness.

Interpretation of our findings is subject to certain limitations. The primary limitation of this study is the cross-sectional study design and our inability to infer the causal relationship between social network variables and loneliness. Second, our data were collected in a rural region of southwestern Uganda and may not generalize to other settings and other populations. Third, although loneliness is reliably measured across many countries, there is insufficient information on the topic in sub-Saharan Africa,



especially in Uganda [63]. Fourth, the measurements of social connections used in this study focused on structural aspects of social connections and did not assess aspects of tie strength, including frequency of contact, reciprocity, and intimacy [64]. As such, additional data are required to provide information about the quality and density of social support to understand the extent to which these may help prevent loneliness among PWH.

## Conclusions

This analysis of cross-sectional data on older PWH and PWOH in rural Uganda identified no statistically significant differences in loneliness, even though PWH had smaller social networks and less financial and physical support. We hypothesize that the quality of social support networks, formal health services, and multiple sources of resilience resources may explain these findings, but further research is needed to better understand the mechanisms underlying the observed differences. While more research is needed to better understand these findings, resolution of this paradox may help to identify interventions for improving the health and mental health of older PWH in rural Uganda.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s10461-023-04258-y>.

**Acknowledgements** We would like to thank the research participants in Uganda and the study team from Mbarara and Kabwohe in Uganda, and the Medical Practice Evaluation Center in the US for making this work possible.

**Author Contributions** All authors contributed to the conception and design of the study. L.T.Q., C.S.R., S.H., Y.T., M.J.S., and A.C.T. contributed to the statistical analyses. L.T.Q., M.J.S., A.C.T., C.S.R., and S.H. contributed to the design of tables and figures. L.T.Q., Z.R., S.A., S.O., E.T., A.F., R.S., and B.O-M contributed to data acquisition. The first draft was written by L.T.Q. All authors contributed to data interpretation, critically reviewed the first draft, and approved the final version.

**Funding** This work was supported by the US National Institutes of Health (R01HL141053, R01AG059504, K43TW010715, K24HL166024, R01MH125667, R01MH113494, and P30AI060354).

## Declarations

**Competing Interests** ACT reports receiving a financial honorarium from Elsevier, Inc. for his work as Co-Editor in Chief of the Elsevier-owned journal *SSM-Mental Health*. The other authors declare no conflicts of interest.

**Sponsor's Role** The findings and conclusions in this report are those of the authors and do not necessarily represent the official views of the US National Institutes of Health.

## References

1. WHO. HIV/AIDS. 2020. [https://www.who.int/health-topics/hiv-aids#tab=tab\\_1](https://www.who.int/health-topics/hiv-aids#tab=tab_1). WHO. Retrieved from [https://www.who.int/health-topics/hiv-aids#tab=tab\\_1](https://www.who.int/health-topics/hiv-aids#tab=tab_1).
2. Greene M, Hessel NA, Perissinotto C, Zepf R, Parrott AH, Foreman C, et al. Loneliness in older adults living with HIV. *AIDS Behav.* 2018;22(5):1475–1484. <https://doi.org/10.1007/s10461-017-1985-1>.
3. Yoo-Jeong M, Brown MJ, Waldrop D. Loneliness mediates the Effect of HIV-related stigma on depressive symptoms among older persons living with HIV. *AIDS Behav.* 2022;26(9):3147–52. <https://doi.org/10.1007/s10461-022-03653-1>.
4. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social isolation as risk factors for mortality: a meta analytic review. *Perspect Psychol Sci.* 2015;10(2):227–37. <https://doi.org/10.1177/1745691614568352>.
5. Crowe CL, Domingue BW, Graf GH, Keyes KM, Kwon D, Bel-sky DW. Associations of loneliness and social isolation with health span and life span in the U.S. *Health and Retirement Study. J Gerontol Ser A.* 2021;76(11):1997–2006. <https://doi.org/10.1093/gerona/glab128>.
6. Cacioppo JT, Hawkley LC, Crawford LE, Ernst JM, Burleson MH, Kowalewski RB, et al. Loneliness and health: potential mechanisms. *Psychosom Med.* 2002;64(3):407–17. <https://doi.org/10.1097/00006842-200205000-00005>.
7. Valtorta N, Hanratty B. Loneliness, isolation and the health of older adults: do we need a new research agenda? *J R Soc Med.* 2012;105(12):518–22. <https://doi.org/10.1258/jrsm.2012.120128>.
8. Harris M, Brouillette M-J, Scott SC, Smaill F, Smith G, Thomas R, et al. Impact of loneliness on brain health and quality of life among adults living with HIV in Canada. *J Acquir Immune Defic Syndr.* 2020;84(4):336. <https://doi.org/10.1097/QAI.0000000000002355>.
9. UN-Habitat. World Cities Report 2016: Urbanization and development - emerging futures. 2016. <https://unhabitat.org/world-cities-report-2016>. Accessed 11 Sept 2023.
10. Payne CF, Pesando LM, Kohler H-P. Private intergenerational transfers, family structure, and health in a sub-saharan African context. *Popul Dev Rev.* 2019;45(1):41–80. <https://doi.org/10.1111/padr.12225>.
11. Hoare E, Jacka F, Berk M. The impact of urbanization on mood disorders: an update of recent evidence. *Curr Opin Psychiatry.* 2019;32(3):198. <https://doi.org/10.1097/YCO.0000000000000487>.
12. Gutu Sakketa T. Urbanisation and rural development in Sub-Saharan Africa: a review of pathways and impacts. *Res Globalization.* 2023;6:100133. <https://doi.org/10.1016/j.resglo.2023.100133>.
13. Barreto M, Victor C, Hammond C, Eccles A, Richins MT, Qualter P. Loneliness around the world: age, gender, and cultural differences in loneliness. *Pers Individ Differ.* 2021;169:110066. <https://doi.org/10.1016/j.paid.2020.110066>.
14. Brennan-Ing M, Kaufman JE, Porter KE, MacPhail C, Seeley J, Karpiak SE, et al. (2022). A detailed analysis of the social support networks of older adults with HIV in Uganda and South Africa. In Brennan-Ing M, Porter KE, Kaufman JE, MacPhail C, Seeley J, editors. *Aging with HIV in Sub-Saharan Africa: Health and Psychosocial Perspectives*. Cham: Springer International Publishing; 2022. p. 139–175. [https://doi.org/10.1007/978-3-030-96368-2\\_8](https://doi.org/10.1007/978-3-030-96368-2_8).
15. Fazeli PL, Moore RC, Vance DE. Resilience attenuates the association between neurocognitive functioning and everyday functioning in individuals aging with HIV in the Deep South. *Int J Geriatr Psychiatry.* 2019;34(1):72–8. <https://doi.org/10.1002/gps.4988>.

16. Kalomo EN, Jun JS, Lee K, Kaddu MN. HIV stigma, resilience and depressive symptoms among older adults living with HIV in rural Namibia. *Afr J AIDS Res.* 2020;19(3):198–205. <https://doi.org/10.2989/16085906.2020.1797841>.
17. Emler CA, Tozay S, Raveis VH. I'm not going to die from the AIDS: resilience in aging with HIV disease. *Gerontologist.* 2011;51(1):101–11. <https://doi.org/10.1093/geront/gnq060>.
18. De Santis JP, Florom-Smith A, Vermeesch A, Barroso S, DeLeon DA. Motivation, management, and mastery: a theory of resilience in the context of HIV infection. *J Am Psychiatr Nurses Assoc.* 2013;19(1):36–46. <https://doi.org/10.1177/1078390312474096>.
19. Cele L, Willen SS, Dhanuka M, Mendenhall E. Ukuphumelela: Flourishing and the pursuit of a good life, and good health, in Soweto, South Africa. *SSM - Mental Health.* 2021;1:100022. <https://doi.org/10.1016/j.ssmmh.2021.100022>.
20. Nyirenda M, Chatterji S, Falkingham J, Mutevedzi P, Hosegood V, Evandrou M, et al. An investigation of factors associated with the health and well-being of HIV-infected or HIV-affected older people in rural South Africa. *BMC Public Health.* 2012;12:259. <https://doi.org/10.1186/1471-2458-12-259>.
21. Negin J, Nyirenda M, Seeley J, Mutevedzi P. Inequality in health status among older adults in Africa: the surprising impact of anti-retroviral treatment. *J Cross-Cult Gerontol.* 2013;28(4):491–3. <https://doi.org/10.1007/s10823-013-9215-4>.
22. Johnson Shen M, Freeman R, Karpiak S, Brennan-Ing M, Seidel L, Siegler EL. The intersectionality of Stigmas among Key populations of older adults affected by HIV: a thematic analysis. *Clin Gerontologist.* 2019;42(2):137–49. <https://doi.org/10.1080/07317115.2018.1456500>.
23. Fakoya OA, McCorry NK, Donnelly M. Loneliness and social isolation interventions for older adults: a scoping review of reviews. *BMC Public Health.* 2020;20(1):129. <https://doi.org/10.1186/s12889-020-8251-6>.
24. Donovan NJ, Blazer D. Social isolation and loneliness in older adults: review and commentary of a National academies Report. *Am J Geriatr Psychiatry.* 2020;28(12):1233–44. <https://doi.org/10.1016/j.jagp.2020.08.005>.
25. Sadler WA. Review of loneliness: the experience of emotional and social isolation. *Contemp Sociol.* 1975;4(2):171–3. <https://doi.org/10.2307/2062224>.
26. Ojembe BU, Kalu ME, Donatus Ezulike C, Iwuagwu AO, Ekoh PC, Oyinlola O, et al. Understanding social and emotional loneliness among black older adults: a scoping review. *J Appl Gerontol.* 2022;41(12):2594–608. <https://doi.org/10.1177/07334648221118357>.
27. Cacioppo JT, Cacioppo S. The phenotype of loneliness. *Eur J Dev Psychol.* 2012;9(4):446–52. <https://doi.org/10.1080/17405629.2012.690510>.
28. van Tilburg TG. Social, emotional, and existential loneliness: a test of the multidimensional concept. *Gerontol.* 2021;61(7):e335–44. <https://doi.org/10.1093/geront/gnaa082>.
29. Weiss RS. Loneliness: The experience of emotional and social isolation. Cambridge, MA, US: The MIT Press. 1973.
30. Russell D, Cutrona CE, Rose J, Yurko K. Social and emotional loneliness: an examination of Weiss's typology of loneliness. *J Personal Soc Psychol.* 1984;46(6):1313–21. <https://doi.org/10.1037//0022-3514.46.6.1313>.
31. Holt-Lunstad J, Robles TF, Sbarra DA. Advancing social connection as a public health priority in the United States. *Am Psychol.* 2017;72(6):517–30. <https://doi.org/10.1037/amp000103>.
32. Quach LT, Ritchie CS, Tsai AC, Reynolds Z, Paul R, Seeley J, et al. The benefits of care: treated HIV infection and health-related quality of life among older-aged people in Uganda. *Aging Ment Health.* 2022:1–7. <https://doi.org/10.1080/13607863.2022.2150143>.
33. Takada S, Nyakato V, Nishi A, O'Malley AJ, Kakuhiire B, Perkins JM, et al. The social network context of HIV stigma: population-based, sociocentric network study in rural Uganda. *Soc Sci Med.* 2019;233:229–36. <https://doi.org/10.1016/j.socscimed.2019.05.012>.
34. Siedner MJ, Bibangambah P, Kim J-H, Lankowski A, Chang JL, Yang IT, et al. Treated HIV infection and progression of carotid atherosclerosis in rural Uganda: a prospective observational cohort study. *J Am Heart Assoc.* 2021;10(12):e019994. <https://doi.org/10.1161/JAHA.120.019994>.
35. Russell DW. UCLA Loneliness Scale (Version 3): reliability, validity, and factor structure. *J Pers Assess.* 1996;66(1):20–40. [https://doi.org/10.1207/s15327752jpa6601\\_2](https://doi.org/10.1207/s15327752jpa6601_2).
36. Steptoe A, Shankar A, Demakakos P, Wardle J. Social isolation, loneliness, and all-cause mortality in older men and women. *Proc Natl Acad Sci USA.* 2013;110(15):5797–801. <https://doi.org/10.1073/pnas.1219686110>.
37. Camlin CS, Charlebois ED, Getahun M, Akatukwasa C, Atwine F, Itiakorit H, et al. Pathways for reduction of HIV-related stigma: a model derived from longitudinal qualitative research in Kenya and Uganda. *J Int AIDS Soc.* 2020;23(12):e25647. <https://doi.org/10.1002/jia2.25647>.
38. Broadhead WE, Gehlbach SH, de Gruy FV, Kaplan BH. The Duke-UNC Functional Social Support Questionnaire. Measurement of social support in family medicine patients. *Med Care.* 1988;26(7):709–23. <https://doi.org/10.1097/00005650-198807000-00006>.
39. Kakuhiire B, Satinsky EN, Baguma C, Rasmussen JD, Perkins JM, Gumisiriza P, et al. Correlates of attendance at community engagement meetings held in advance of bio-behavioral research studies: A longitudinal, sociocentric social network study in rural Uganda. *PLoS Med.* 2021;18(7):e1003705. <https://doi.org/10.1371/journal.pmed.1003705>.
40. SAS: Analytics, Artificial Intelligence and Data Management. (2013, July). Retrieved June 23, 2022, from [https://www.sas.com/en\\_us/home.html](https://www.sas.com/en_us/home.html).
41. Schafer MH, Upenieks L, DeMaria J. Do older adults with HIV have distinctive personal networks? Stigma, network activation, and the role of disclosure in South Africa. *AIDS Behav.* 2021;25(5):1560–72. <https://doi.org/10.1007/s10461-020-02996-x>.
42. Reynolds Z, Gilbert R, Sentongo R, Meyer A-C, Saylor D, Okello S, et al. (2022). Priorities for health and wellbeing for older people with and without HIV in Uganda: a qualitative methods study. *J Int AIDS Soc.* 2022;25 Suppl 4:e26000. <https://doi.org/10.1002/jia2.26000>.
43. Dulin AJ, Dale SK, Earnshaw VA, Fava JL, Mugavero MJ, Napravnik S, et al. (2018). Resilience and HIV: A review of the definition and study of resilience. *AIDS Care* 2018;30 Suppl 5:S6–S17. <https://doi.org/10.1080/09540121.2018.1515470>.
44. Woodward EN, Banks RJ, Marks AK, Pantalone DW. Identifying Resilience resources for HIV Prevention among sexual minority men: a systematic review. *AIDS Behav.* 2017;21(10):2860–73. <https://doi.org/10.1007/s10461-016-1608-2>.
45. Whitson HE, Duan-Porter W, Schmader KE, Morey MC, Cohen HJ, Colón-Emeric CS. Physical resilience in older adults: systematic review and development of an emerging construct. *The J Gerontol Ser A.* 2016;71(4):489–95. <https://doi.org/10.1093/gerona/glv202>.
46. De Jesus M, Ware D, Brown AL, Egan JE, Haberlen SA, Paella FJ, et al. Social-environmental resiliencies protect against loneliness among HIV-positive and HIV-negative older men who have sex with men: results from the multicenter aids cohort study (MACS). *Soc Sci Med.* 2021;272:113711. <https://doi.org/10.1016/j.socscimed.2021.113711>.




47. Brennan-Ing M, Porter KE, Kaufman JE, MacPhail C, Seeley J, editors. *Aging with HIV in Sub-Saharan Africa: health and psychosocial perspectives*. Cham: Springer International Publishing; 2022. <https://doi.org/10.1007/978-3-030-96368-2>.
48. Mburu G, Ram M, Skovdal M, Bitira D, Hodgson I, Mwai GW, et al. Resisting and challenging stigma in Uganda: the role of support groups of people living with HIV. *J Int AIDS Soc*. 2013;16(3 Suppl 2):18636. <https://doi.org/10.7448/IAS.16.3.18636>.
49. Geldsetzer P, Vaikath M, Wagner R, Rohr JK, Montana L, Gómez-Olivé FX, et al. Depressive symptoms and their relation to age and chronic diseases among middle-aged and older adults in rural South Africa. *J Gerontol - Biol Sci Med Sci*. 2019;74(6):957–63. <https://doi.org/10.1093/gerona/gly145>.
50. Tsai AC, Bangsberg DR, Bwana M, Haberer JE, Frongillo EA, Muzoora C, et al. (2013). How does antiretroviral treatment attenuate the stigma of HIV? Evidence from a cohort study in rural Uganda. *AIDS Behav*. 2013;17(8):2725–31. <https://doi.org/10.1007/s10461-013-0503-3>.
51. Wandera SO, Ddumba I, Akinyemi JO, Adedini SA, Odimegwu C. Living alone among older persons in Uganda: prevalence and associated factors. *Ageing Int*. 2017;42(4):429–46. <https://doi.org/10.1007/s12126-017-9305-7>.
52. Chang LW, Nakigozi G, Billioux VG, Gray RH, Serwadda D, Quinn TC, et al. (2015). Effectiveness of peer support on care engagement and preventive care intervention utilization among pre-antiretroviral therapy, HIV-infected adults in Rakai, Uganda: a randomized trial. *AIDS Behav*. 2015;19(10):1742–1751. <https://doi.org/10.1007/s10461-015-1159-y>.
53. Schatz E, Seeley J. Gender, ageing & carework in East and Southern Africa: a review. *Glob Public Health*. 2015;10(10):1185–200. <https://doi.org/10.1080/17441692.2015.1035664>.
54. Cooper F, Lewis EG, Urasa S, Whitton L, Collin H, Coles S, et al. Social vulnerability, frailty, and their association with mortality in older adults living in rural Tanzania. *J Gerontol A Biol Sci Med Sci*. 2022;77(10):2050–58. <https://doi.org/10.1093/gerona/glac066>.
55. Stephens C, Alpass F, Towers A, Stevenson B. The effects of types of social networks, perceived social support, and loneliness on the health of older people: accounting for the social context. *J Aging Health*. 2011;23(6):887–911. <https://doi.org/10.1177/0898264311400189>.
56. Geffen LN, Kelly G, Morris JN, Howard EP. Peer-to-peer support model to improve quality of life among highly vulnerable, low-income older adults in Cape Town, South Africa. *BMC Geriatr*. 2019;19(1):279. <https://doi.org/10.1186/s12877-019-1310-0>.
57. Hao G, Bishwajit G, Tang S, Nie C, Ji L, Huang R. Social participation and perceived depression among elderly population in South Africa. *Clin Interv Aging*. 2017;12:971–6. <https://doi.org/10.2147/CIA.S137993>.
58. Goveas JS, Ray RM, Woods NF, Manson JE, Kroenke CH, Michael YL, et al. Associations between changes in loneliness and social connections, and mental health during the COVID-19 pandemic: the women's health initiative. *J Gerontol A Biol Sci Med Sci*. 2022;77 Suppl 1:S31–S41. <https://doi.org/10.1093/gerona/glab371>.
59. Boehlen FH, Maatouk I, Friederich H-C, Schoettker B, Brenner H, Wild B. Loneliness as a gender-specific predictor of physical and mental health-related quality of life in older adults. *Qual Life Res*. 2022;31(7):2023–33. <https://doi.org/10.1007/s11136-021-03055-1>.
60. Pagan R. Gender and age differences in loneliness: evidence for people without and with disabilities. *Int J Environ Res Public Health*. 2020;17(24):9176. <https://doi.org/10.3390/ijerph17249176>.
61. Girgus JS, Yang K, Ferri CV. The gender difference in depression: are elderly women at greater risk for depression than elderly men? *Geriatrics*. 2017;2(4):35. <https://doi.org/10.3390/geriatrics2040035>.
62. Kim YB, Lee SH. Gender differences in correlates of loneliness among community-dwelling older Koreans. *Int J Environ Res Public Health*. 2022;19(12):7334. <https://doi.org/10.3390/ijerph19127334>.
63. Surkalim DL, Luo M, Eres R, Gebel K, van Buskirk J, Bauman A, Ding D. The prevalence of loneliness across 113 countries: systematic review and meta-analysis. *BMJ*. 2022;e067068. <https://doi.org/10.1136/bmj-2021-067068>.
64. Tsai AC, Papachristos AV. From social networks to health: Durkheim after the turn of the millennium. Introduction. *Soc Sci Med*. 2015;125:1–7. <https://doi.org/10.1016/j.socscimed.2014.10.045>.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

## Authors and Affiliations

Lien T. Quach<sup>1,2,3,4,5,16</sup>  · Christine S. Ritchie<sup>2,3,5</sup> · Zahra Reynolds<sup>1</sup> · Robert Paul<sup>6</sup> · Janet Seeley<sup>7</sup> · Yao Tong<sup>1</sup> · Susanne Hoepfner<sup>5,8</sup> · Samson Okello<sup>9,10</sup> · Noeline Nakasujja<sup>11</sup> · Brianne Olivieri-Mui<sup>12</sup> · Deanna Saylor<sup>13</sup> · Meredith Greene<sup>14</sup> · Stephen Asimwe<sup>1,9,15</sup> · Edna Tindimwebwa<sup>15</sup> · Flavia Atwiine<sup>9</sup> · Ruth Sentongo<sup>9</sup> · Mark J. Siedner<sup>1,5,9</sup> · Alexander C. Tsai<sup>5,8,9</sup>

✉ Lien T. Quach  
lquach@mgh.harvard.edu

<sup>1</sup> Medical Practice Evaluation Center, Mongan Institute, Department of Medicine, Massachusetts General Hospital, Boston, MA, USA

<sup>2</sup> Centre for Aging and Serious Illness, Mongan Institute, Massachusetts General Hospital, Boston, MA, USA

<sup>3</sup> Division of Palliative Care and Geriatric Medicine, Department of Medicine, Massachusetts General Hospital, Boston, USA

<sup>4</sup> Department of Gerontology, University of Massachusetts Boston, Boston, MA, USA

<sup>5</sup> Harvard Medical School, Boston, MA, USA

<sup>6</sup> Department of Psychological Sciences, University of Missouri at St. Louis, St. Louis, USA

<sup>7</sup> Department of Global Health and Development, Faculty of Public Health and Policy, London School of Hygiene & Tropical Medicine, London, UK

<sup>8</sup> Department of Psychiatry, Massachusetts General Hospital, Boston, MA, USA

<sup>9</sup> Mbarara University of Science and Technology, Mbarara, Uganda

<sup>10</sup> Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina, Chapel Hill, NC, USA

<sup>11</sup> Department of Psychiatry, College of Health Sciences, Makerere University School of Medicine, Kampala, Uganda

<sup>12</sup> Department of Health Sciences, Bouvé College of Health Sciences, Northeastern University, Boston, MA, USA

<sup>13</sup> Department of Neurology, Johns Hopkins University School of Medicine, Baltimore, MD, USA

<sup>14</sup> Department of Medicine, Indiana University School of Medicine and Indiana University Center for Aging Research, Regenstreif Institute, Inc., Indianapolis, IN, USA

<sup>15</sup> Kabwohe Clinical Research Center, Kabwohe, Uganda

<sup>16</sup> The Massachusetts Veterans Epidemiology Research and Information Center, VA Boston Healthcare System, Boston, USA